

APPLICATIONS OF AUTOMATA AND GRAPHS: LABELING-OPERATORS IN HILBERT SPACE II

ILWOO CHO AND PALLE E. T. JORGENSEN

ABSTRACT. We introduced a family of infinite graphs directly associated with a class of von Neumann automaton model $\mathcal{A}_G$. These are finite state models used in symbolic dynamics: stimuli models and in control theory. In the context of groupoid von Neumann algebras, and an associated fractal group, we prove a classification theorem for representations of automata.

Table of Contents

1. Introduction	2
2. Background	7
2.1. Amalgamated Free Probability	7
2.2. Basic Concepts	9
3. Definitions	12
3.1. Categorical Groupoids & Groupoid Actions	12
3.2. Automata and Fractal Groups	14
3.3. Right Graph von Neumann Algebras	18
3.4. M -Valued Right Graph W^* -Probability Spaces	21
4. Labeled Graph Groupoids and Graph Automata	26
4.1. Labeled Graph Groupoids	27
4.2. The Operation θ and ω_+	34
4.3. Graph Automata	35
4.4. Graph-Automata Trees	38
5. Labeling Operators of Graph Groupoids	43
5.1. The Labeling Operator of $\mathbb{G}$ in M_G	43
5.2. D_G -Valued Free Distributional Data of Labeling Operators	45
5.3. Graph-Theoretical Characterization of $\mathcal{W}_{2n}^{(m)}$ and $\mathcal{W}_{2n}^{(c)}$	52
6. Examples	54
References	57

Date: Mar., 2008.

1991 *Mathematics Subject Classification.* 05C62, 05C90, 17A50, 18B40, 46K10, 47A67, 47A99, 47B99.

Key words and phrases. Graph Groupoids, Labeled Graph Groupoids, Graph Automata, Graph-Groupoid-Automata, Automata-Trees, Right Graph von Neumann Algebras, Right Graph W^* -Probability Spaces, Labeling Operators.

The second named author was supported by the U. S. National Science Foundation.

1. INTRODUCTION

This paper is the second of two papers studying representations of graphs by operators in Hilbert space, and their applications, with the help of Automata Theory. While graph theory is traditionally considered part of discrete mathematics, in this paper we show that applications of tools from automata and operators on Hilbert spaces yield global results for representations of a class of infinite graphs, as well as spin-off applications. We begin with an outline of the use of automata, and more generally, of finite state models (FSMs) in the processing of numbers, or more importantly in sampling and in quantization of digitized information such as speech signals and digital images. In these models, the finite input states of a particular FSM might be frequency-bands (for example a prescribed pair of high-pass and low-pass digital filters), or a choice of subdivision filters; where the subdivision refers to data sets with self-similarity; such as is typically seen in fractals. Hence, these applications make connections to discrete wavelet algorithms as used in signal and image processing, as well as in science and engineering. If the input-options for a particular FSM are chosen from a prescribed system of low-pass and high-pass filters, the resulting discrete model can then be realized by operators on Hilbert spaces. Similarly, images are digitized into matrix shapes used in computer programs for compression of images.

In a general operator theoretic setting, this paper introduces the relevant representations of the generators of graphs and automata. Hence data from FSMs and graphs are represented with the use of Hilbert space geometry. Recall that digital images are typically given by matrices of pixels, and that spectral analysis and matrix operations can be done with operators in Hilbert space. However, the Hilbert spaces needed in a particular application are typically not immediately apparent from the particular engineering problem under discussion. This paper focused on making the link between graphs and automata on the discrete side to Hilbert space operators and representations on the spectral side.

The list of the advantages deriving from the use of operator algebraic methods in the analysis of graphs includes the following: starting with a particular graph G , a main question is that of identifying decompositions and global invariants for G . Motivated by ergodic theory, a further goal is to analyze global properties of G in terms of its local features, and an operator algebraic decomposition is a powerful tool in the identification of local vs global features and invariants of G . For the purpose, one may use that the theory of von Neumann algebras (rings of operators) allows factor decompositions (See [66]). And the von Neumann factors then further allow type classification, type I , II , and III , with sub-types; See [66] for the fundamentals in the theory of von Neumann algebras, and [64], [65] for some relevant applications to infinite graphs of this idea. In addition, the analysis of graphs uses infinite determinant, and the Fuglede-Kadison determinant [63] (from von Neumann algebra theory) is a powerful tool in graph computation because it passes to direct Hilbert space decompositions.

The potential-theoretic part of the subject may be understood as a mathematical idealization of electrical networks on infinite graphs G (See [43], [44] and [45]). We address two issues from Analysis: Find representations (by operators on Hilbert spaces) of the graph systems G , and identify a class of operators whose spectral theory captures significant information about G . We focus on the graphs themselves,

and our motivation derives in part from work by Strichartz and Kigami (See [46], [47] and [48]) and others for restricted classes of fractals. The focus there is the adaptation of a rescaling and energy-renormalization on graphs and an adaptation to fractal models X ; for example, to Brownian motion on X , or to a version of differential operators on spaces of functions on X . Here, our focus is on Operator Theory needed for analysis of graphs and automata (finite-state machines: e.g., [54]) such as are used in for example signal processing algorithm, e.g., those based on a discrete multiresolution (e.g., see [52]).

We begin our paper with an outline of three trends: (i) Hilbert space and Spectral Analysis on the graphs, (ii) Analysis on associated fractals arising from automata, and finally (iii) the interplay between (i) and (ii). We further stress some of the differences between the two.

The main purpose of this paper is to introduce a new algebraic structures having certain fractal property. In [10], [11], [13], [14] and [15], we introduced graph groupoids induced by countable directed graphs. A graph groupoid is a categorial groupoid having as a base the set of all vertices of the given graph. We know that every groupoid having only one base element is a group. So, if G is a finite directed graph with graph groupoid $\mathbb{G}$, and if the vertex set $V(G) \subset \mathbb{G}$ consists of only one element, then the graph groupoid $\mathbb{G}$ of this one-vertex- n -loop-edge graph G , for $n = |E(G)| \in \mathbb{N}$, is a group; furthermore, it is group-isomorphic to the free group F_n with n -generators (See [10] and [11]). Here, $E(G)$ denotes the edge set of G .

Then we can embed the edge set $E(G)$ into the edge set $E(\widehat{G})$, where $\widehat{G}$ is the shadowed graph of G . Put the lattice weights $l_{\pm 1}, \dots, l_{\pm n}$, associate to the edges $e_1^{\pm 1}, \dots, e_n^{\pm 1}$ in $E(\widehat{G})$, where $e_j^{-1} \in E(\widehat{G})$ means the shadow of $e_j \in E(G)$ (See Section 2), where $l_{\pm j} = (0, \pm e^j)$ in $\mathbb{R}^2$. Here, e is the natural number in $\mathbb{R}$. More precisely, we define the labeling set $\mathcal{X}_0$ by $\pm X = \{l_{\pm 1}, \dots, l_{\pm n}\}$. Then define the labeling map

$$\varphi : \mathcal{X}_0 \times E(\widehat{G}) \rightarrow \mathcal{X}_0$$

by

$$\varphi(l, e) = \text{the weight } l_e \text{ of } e.$$

Also, define the shifting map

$$\psi : \mathcal{X}_0 \times E(\widehat{G}) \rightarrow E(\widehat{G})$$

by

$$\psi(l, e) = e.$$

Then we have the automaton $\mathcal{A}_G = \langle \pm X_0, E(\widehat{G}), \varphi, \psi \rangle$. Then this automaton $\mathcal{A}_G = \langle \pm X_0, E(\widehat{G}), \varphi, \psi \rangle$ generates a fractal group $\Gamma \stackrel{\text{Group}}{=} F_n$. We further address the general case where the vertex set contains more than one element? The answer of this question was provided in [17]. In this paper, we will consider more general case than this.

Our paper is organized as follows. The remaining of Introduction contains an overview, including some definitions; meant only as sketch. Full definitions and statements of theorems will be in the respective main sections of the paper.

Section 2 introduces those fundamentals from amalgamated free probability theory which will be needed later. Here, we use the term “free probability” in the sense of D. Voiculescu (e.g., See [5]), a framework where random variables are noncommutative operators in Hilbert space, as opposed to commuting measurable functions on a probability space. While Voiculescu’s notion free independence is modeled on the classical concept of independence of families of random variables, there are important differences. They are reflected in the other notions we need from the noncommutative setting: operator-valued $*$ -moments and $*$ -cumulants. In addition, we will need fundamentals from the theory of partitions. This material allows applications of operator algebra tools to our present graph analysis of automata. Moreover, within this framework, we will need an extension of the notion of group crossed products from operator algebra theory: we introduce a (groupoid) crossed product of a von Neumann algebra by a groupoid (See Definition 2.1).

Section 3 focuses on graphs and graph groupoids: categorial groupoids and groupoid actions. Here, actions refers to actions by transformations on von Neumann algebras.

A main application will be to von Neumann’s automata (i.e., certain self-reproducing dynamical structures) introduced in Section 3.2. From each automaton $\mathcal{A}$, we define an automata group $\Gamma(\mathcal{A})$, and we identify a subclass of $\mathcal{A}$ for which the corresponding group is a fractal (Definition 3.1). The rest of Section 3 deals with actions on von Neumann algebras which are induced by countable directed graphs. Starting with the graph, in Theorem 3.2, we show that a groupoid crossed product von Neumann algebra $\mathbb{M}_G$ has an amalgamated free structures over its diagonal W^* -subalgebra $\mathbb{D}_G$. We then use this in Theorem 3.4 and 3.5 in breaking up a given W^* -probability space into basic constituents. Specially, we show that every groupoid crossed product algebra $\mathbb{M}_G$, as a W^* -probability space, decomposes in a specific way into a direct decomposition in such a manner that the constituents in the decomposition are so-called $\mathbb{D}_G$ -free blocks. Moreover, Theorem 3.6 offers a classification of the $\mathbb{D}_G$ -free blocks which may occur. Applications, Example 3.5 and 3.6 are given to edge graphs, while Section 4 covers graph-automata (Definition 4.7). In our further analysis of automata-labeled graph groupoids and their von Neumann algebras, this is brought to bear. In Section 4.4, we specialize to graph-automata trees, spelling out more detailed results.

The theorems from Section 3 and 5 are strengthened in the remaining two sections of our paper with the use of labeling operators. Theorem 5.4 makes use of a condition for labeling operators in deciding $\mathbb{D}_G$ -freeness, while Theorem 5.6 and 5.8 yield explicit formulas for $\mathbb{D}_G$ -valued joint moments, and joint cumulants, respectively. This is then further refined, and applied in Section 6, the fractal case in Example 6.1.

We realize that, by giving weights on the edges of a given graph G , which are determined by the out-degrees of vertices, we can get the corresponding automaton $\mathcal{A}_G$. By observing the properties of $\mathcal{A}_G$, we can determine the groupoidal version of a fractal property, relative to a fractal group. We will say that a graph groupoid $\mathbb{G}$ is a fractaloid, if $\mathcal{A}_G$ acts “fully” on a $|V(\widehat{G})|$ -copies of regular growing directed trees. For example, if a directed graph G is a one-flow circulant graph, then the graph groupoid $\mathbb{G}$ is a fractaloid. How about the given graph groupoids are not fractaloids? This is our motivation of this work.

Similar to [10] and like [17], we can have a right graph von Neumann algebra $\mathbb{M}_G = \overline{\mathbb{C}[\beta(\mathbb{G})]}^w$, as a W^* -subalgebra of the operator algebra $B(H_G)$, consisting of all bounded linear operators on H_G , where H_G is a generalized Fock space, called the graph Hilbert space induced by $\mathbb{G}$. We are interested in the case where $\mathbb{G}$ is a fractaloid. Then, similar to the classical case, we can define the Hecke-type operator $\tau \in \mathbb{M}_G$. Instead of considering the pure operator-theoretical data of τ , we will observe the amalgamated free distributional data of τ . Since τ is self-adjoint, the amalgamated free moments of it contains the operator-valued spectral measure theoretical information. This means that the free moments of τ will contain the operator-theoretical data of τ . In [17], we observed the special cases where the given graph groupoids are fractaloids.

From the theory of algebras of operators in Hilbert space, we will need von Neumann algebra constructions (e.g., [59]), free probability, in particular, amalgamated free products (e.g., [5] and [23]), groupoids and groupoid actions (e.g., [19] and [60]), and Hecke operators (e.g., [27]). In technical discussions, we will use “von Neumann algebras” and “ W^* -algebras” synonymously. If H is a given Hilbert space, the algebra of all bounded operators on H is denoted by $B(H)$. From Graph Theory, we use such notations as “sets of vertices”, “sets of edges”, “loops” and “degrees”, etc. If G is a given directed graph, we introduce a “shadow” construction from a reversal of edges, denoted by G^{-1} ; see details below. From symbolic dynamics, we shall use fundamentals of “automata”, as well as free constructions, such as the free group with multi-generators. Hence, our next section will contain a number of definitions that will be needed later.

Since our paper is interdisciplinary and directed at several audiences, we have included details from one area of mathematics which might not be familiar to readers from another.

A *graph* is a set of objects called *vertices* (or points or nodes) connected by links called *edges* (or lines). In a *directed graph*, the two directions are counted as being distinct directed edges (or arcs). A graph is depicted in a diagrammatic form as a set of dots (for vertices), jointed by curves (for edges). Similarly, a directed graph is depicted in a diagrammatic form as a set of dots jointed by arrowed curves, where the arrows point the direction of the directed edges.

Throughout this paper, every graph is a locally finite countably directed graph. Recall that we say that a countably directed graph G is *locally finite* if each vertex of G has only finitely many incident edges. Equivalently, the degree of v is finitely

determined. Also, recall that the degree $\deg(v)$ of a vertex v is defined to be the sum of the out-degree $\deg_{out}(v)$ and the in-degree $\deg_{in}(v)$, where

$$\deg_{out}(v) \stackrel{def}{=} |\{e \in E(G) : e \text{ has its initial vertex } v\}|$$

and

$$\deg_{in}(v) \stackrel{def}{=} |\{e \in E(G) : e \text{ has its terminal vertex } v\}|.$$

Let $\widehat{G}$ be the shadowed graph of G , in the sense of [10]. Then we can consider the degree of each vertex of $\widehat{G}$, too, since $\widehat{G}$ is also a locally finite countable directed graph. Assume that

$$N = \max \{\deg_{out}(v) : v \in V(\widehat{G}) = V(G)\}.$$

Then we can define the labeling set $X = \{1, 2, \dots, N\}$. We assign the weights $\{1, \dots, N\}$ to all elements in the edge set $E(\widehat{G})$ of the shadowed graph $\widehat{G}$ of G . (This weighting provides the weights of all elements in $\mathbb{G}$, which are the sequences contained in X_0^∞ , where $X_0 = \{0\} \cup X$.) We will call this process placing the weights onto all elements of $\mathbb{G}$, the labeling process. This labeling process lets us construct the automaton $\mathcal{A}_G = \langle X_0, \mathbb{G}, \varphi, \psi \rangle$ induced by the graph groupoid $\mathbb{G}$. If the automaton $\mathcal{A}_G$ satisfies certain *fractal* property; we will call the graph groupoid $\mathbb{G}$ a fractaloid. Clearly, the word “fractaloid” hints at “fractal (graph) groupoid”.

As in [10] and [11], we will fix a representation (H_G, β) of a graph groupoid $\mathbb{G}$, where H_G is the graph Hilbert space and β is a certain groupoid action of $\mathbb{G}$. Let $X = \{x_1, x_2, \dots, x_N\}$ be the labeling set. Then we can define the operator $\tau_j \in B(H_G)$ by

$$\tau_j(\xi_w) = \xi_w \xi_e = \xi_{we}, \text{ for all } \xi_w \in \mathcal{B}_{H_G},$$

for each $j \in X$, where $\mathcal{B}_{H_G}$ is the Hilbert basis of H_G , whenever an edge e has its weight x_j , and $we \neq \emptyset$ in $\mathbb{G}$. Then we can have the operator $\tau \in B(H_G)$ defined by

$$\tau = \sum_{j=1}^N \tau_j.$$

This Hecke-type operator on H_G is said to be the labeling operator of $\mathbb{G}$ on H_G . We will consider the free distributional data of this operator τ on H_G . If $\mathbb{G}$ is a fractaloid, then τ is self-adjoint. So, the free moments of it contain the spectral theoretical properties of τ . In particular, we can show that the amalgamated free moments $(E(\tau^n))_{n=1}^\infty$ of τ are determined by the cardinalities $(\eta_n)_{n=1}^\infty$ of certain subsets $\mathcal{W}_n^{(m)}$ of the collection of all finite paths in the shadowed graph $\widehat{G}$ of the given graph G , determined recursively for $n \in \mathbb{N}$. Also, the equivalent free distributional data is provided.

2. BACKGROUND

Recently, the countable directed graphs have been studied in Pure and Applied Mathematics, because not only that they are involved by a certain noncommutative structures but also that they visualize such structures. Furthermore, the visualization has nice matricial expressions, (sometimes, the operator-valued matricial expressions dependent on) adjacency matrices or incidence matrices of the given graph. In particular, partial isometries in an operator algebra can be expressed and visualized by directed graphs: in [10], [11], [13] and [14], we showed that each edge (resp. each vertex) of a graph matches to a partial isometry (resp. a projection) in a (graph) Hilbert space, and in [15], we showed that any partial isometry (resp. projections induced by this partial isometry) on an arbitrary separable infinite Hilbert space matches to an edge (resp. vertices) of certain (corresponding) directed graph.

The main purpose of this paper is to introduce algebraic and operator-algebraic structures induced by countable directed graphs. Basically, we will follow the same settings used in [13]. For convenience, as we assumed at the beginning of this paper, we will restrict our interests to the case where the countable directed graphs are locally finite. In [10] and [11], starting with a countable directed graph, we assign certain algebraic elements gotten from the admissibility, and then we assign partial isometries to those elements. From these operators, we generated a von Neumann algebra and then we considered free probabilistic properties of them. In conclusion, we found the nice amalgamated free block structures of such von Neumann algebras and this provides the full characterization of the von Neumann algebras generated by graph groupoids.

2.1. Amalgamated Free Probability.

Let $B \subset A$ be von Neumann algebras with $1_B = 1_A$ and assume that there is a *conditional expectation* $E_B : A \rightarrow B$ satisfying that (i) E_B is a ($\mathbb{C}$ -)linear map, (ii) $E_B(b) = b$, for all $b \in B$, (iii) $E_B(b_1 a b_2) = b_1 E_B(a) b_2$, for all $b_1, b_2 \in B$ and $a \in A$, (iv) E_B is continuous under the given topologies for A (and B), and (v) $E_B(a^*) = E_B(a)^*$ in B , for all $a \in A$. The algebraic pair (A, E_B) is said to be a *B-valued W^* -probability space*. Every operator in (A, E_B) is called a *B-valued (free) random variable*. Any *B-valued* random variables have their *B-valued* free distributional data: *B-valued* $*$ -moments and *B-valued* $*$ -cumulants of them. Suppose $a_1, \dots, a_s$ are *B-valued* random variables in (A, E_B) , where $s \in \mathbb{N}$. The $(i_1, \dots, i_n)$ -th joint *B-valued* $*$ -moments of $a_1, \dots, a_s$ are defined by

$$E_B \left((b_1 a_{i_1}^{r_{i_1}}) (b_2 a_{i_2}^{r_{i_2}}) \dots (b_n a_{i_n}^{r_{i_n}}) \right),$$

and the $(j_1, \dots, j_k)$ -th joint *B-valued* $*$ -cumulants of $a_1, \dots, a_s$ are defined by

$$k_k^B \left((b_1 a_{j_1}^{r_{i_1}}), \dots, (b_k a_{j_k}^{r_{i_k}}) \right) = \sum_{\pi \in NC(k)} E_{B:\pi} \left(b_1 a_{j_1}^{r_{i_1}}, \dots, b_k a_{j_k}^{r_{i_k}} \right) \mu(\pi, 1_k),$$

for all $(i_1, \dots, i_n) \in \{1, \dots, s\}^n$ and for all $(j_1, \dots, j_k) \in \{1, \dots, s\}^k$, for $n, k \in \mathbb{N}$, where $b_j \in B$ are arbitrary and $r_{i_1}, \dots, r_{i_n}, r_{j_1}, \dots, r_{j_k} \in \{1, *\}$ and $NC(k)$ is the lattice of all *noncrossing partitions* with its minimal element $0_k = \{(1), (2), \dots, (k)\}$ and its maximal element $1_k = \{(1, 2, \dots, k)\}$, for all $k \in \mathbb{N}$, and μ is the *Moebius functional* in the incidence algebra $\mathcal{I}$. Here, $E_{B:\pi}(\dots)$ is the *partition-depending B-valued moment*. For example, if $\pi = \{(1, 4), (2, 3), (5)\}$ in $NC(5)$, then

$$E_{B:\pi}(a_1, a_2, a_3, a_4, a_5) = E_B(a_1 E_B(a_2 a_3) a_4) E_B(a_5).$$

Recall that the lattice $NC(n)$ of all noncrossing partitions over $\{1, \dots, n\}$ has its partial ordering “ $\leq$ ”,

$$\pi \leq \theta \stackrel{\text{def}}{\iff} \text{for each block } V \text{ in } \pi, \exists \text{ a block } B \text{ in } \theta \text{ s.t., } V \subseteq B,$$

for $\pi, \theta \in NC(n)$, where “ $\subseteq$ ” means the usual set inclusion, for all $n \in \mathbb{N}$. Also recall that the *incidence algebra* $\mathcal{I}$ is the collection of all functionals

$$\xi : \cup_{n=1}^{\infty} (NC(n) \times NC(n)) \rightarrow \mathbb{C}$$

satisfying that $\xi(\pi, \theta) = 0$, whenever $\pi > \theta$, with the usual function addition (+) and the *convolution* (*) defined by

$$(\xi_1 * \xi_2)(\pi, \theta) \stackrel{\text{def}}{=} \sum_{\pi \leq \sigma \leq \theta} \xi_1(\pi, \sigma) \xi_2(\sigma, \theta),$$

for all $\xi_1, \xi_2 \in \mathcal{I}$. If we define the *zeta functional* $\zeta \in \mathcal{I}$ by

$$\zeta(\pi, \theta) = 1, \text{ for all } \pi \leq \theta \text{ in } NC(n), \text{ for all } n \in \mathbb{N},$$

then it is the identity element of $\mathcal{I}$ having its convolution inverse, the Moebius functional μ . Thus the Moebius functional μ satisfies that

$$\mu(0_n, 1_n) = (-1)^{n-1} c_{n-1} \text{ and } \sum_{\pi \in NC(n)} \mu(\pi, 1_n) = 0,$$

where $c_k \stackrel{\text{def}}{=} \frac{1}{k+1} \binom{2k}{k}$ is the k -th Catalan number, for all $k \in \mathbb{N}$ (See [21]).

The B -valued freeness on (A, E_B) is characterized by the B -valued $*$ -cumulants. Let A_1 and A_2 be W^* -subalgebras of A having their common W^* -subalgebra B . We say that A_1 and A_2 are *free over B* in (A, E_B) if all mixed B -valued $*$ -cumulants

of A_1 and A_2 vanish. The subsets X_1 and X_2 of A are said to be free over B in (A, E_B) if the W^* -subalgebras $vN(X_1, B)$ and $vN(X_2, B)$ of A are free over B in (A, E_B) , where $vN(S_1, S_2)$ means the von Neumann algebra generated by sets S_1 and S_2 . Similarly, we say that the B -valued random variables x and y are free over B in (A, E_B) if the subsets $\{x\}$ and $\{y\}$ are free over B in (A, E_B) (Also See [21]).

Suppose two W^* -subalgebras A_1 and A_2 of A , containing their common W^* -subalgebra B , are free over B in (A, E_B) . Then we can create the W^* -subalgebra $vN(A_1, A_2)$ of A generated by A_1 and A_2 . It is denoted by $A_1 *_B A_2$, which is called the B -valued free product (sub)algebra of A_1 and A_2 (over B). Assume that the given von Neumann algebra A is generated by its W^* -subalgebras A_i , containing their common W^* -subalgebra B , for $i \in \Lambda$. Also, assume that all A_i 's are free over B from each other in (A, E_B) . Then A is denoted by $*_{i \in \Lambda} A_i$. i.e., the von Neumann algebra A is said to be a B -valued free product algebra of A_i 's (over B).

Suppose a von Neumann algebra A is a B -free product algebra $*_{i \in \Lambda} A_i$. Then A is Banach-space isomorphic to the Banach space

$$B \oplus \left(\bigoplus_{n=1}^{\infty} \left(\bigoplus_{i_1 \neq i_2, i_2 \neq i_3, \dots, i_{n-1} \neq i_n} (A_{i_1}^o \otimes_B \dots \otimes_B A_{i_n}^o) \right) \right)$$

with

$$A_{i_j}^o \stackrel{def}{=} A_{i_j} \ominus B, \text{ for all } j = 1, \dots, n,$$

where $\otimes_B$ is the B -valued tensor product.

2.2. Basic Concepts.

For a given countable directed graph G , we can define the oppositely directed graph G^{-1} , with $V(G^{-1}) = V(G)$ and $E(G^{-1}) = \{e^{-1} : e \in E(G)\}$, where each element e^{-1} satisfies that $e = v_1 e v_2$ in $E(G)$, with $v_1, v_2 \in V(G)$, if and only if $e^{-1} = v_2 e^{-1} v_1$, in $E(G^{-1})$. This opposite directed edge $e^{-1} \in E(G^{-1})$ of $e \in E(G)$ is called the shadow of e . Also, this new graph G^{-1} , induced by G , is said to be the shadow of G . It is clear that $(G^{-1})^{-1} = G$.

Define the *shadowed graph* $\widehat{G} = G \cup G^{-1}$ of G by a directed graph with its vertex set $V(\widehat{G}) = V(G) = V(G^{-1})$ and its edge set $E(\widehat{G}) = E(G) \cup E(G^{-1})$, where G^{-1} is the *shadow* of G . Then we can construct the *free semigroupoid* $\mathbb{F}^+(\widehat{G})$ of the shadowed graph $\widehat{G}$, as a collection of all vertices and finite paths of $\widehat{G}$ with its binary operation called the *admissibility*, where the admissibility is nothing but the direction-depending connectedness of elements in $\mathbb{F}^+(\widehat{G})$. Notice that all finite paths in $\mathbb{F}^+(\widehat{G})$ are the words in $V(\widehat{G}) \cup E(\widehat{G})$. By defining the *reduction* (RR) on $\mathbb{F}^+(\widehat{G})$, we can construct the graph groupoid $\mathbb{G}$. i.e., the *graph groupoid* $\mathbb{G}$ is a set of all “reduced” words in $E(\widehat{G})$, with the inherited admissibility on $\mathbb{F}^+(\widehat{G})$, where the reduction (RR) on $\mathbb{G}$ is

$$(RR) \quad w w^{-1} = v \text{ and } w^{-1}w = v',$$

for all $w = v w v' \in \mathbb{G}$, with $v, v' \in V(\widehat{G})$. In fact, this graph groupoid $\mathbb{G}$ is indeed a categorial groupoid with its base $V(\widehat{G})$.

For an arbitrary fixed von Neumann algebra M in an operator algebra $B(K)$ of all bounded linear operators on a Hilbert space K , we define a crossed product algebra $\mathbb{M}_G = M \times_\beta \mathbb{G}$ of M and $\mathbb{G}$ via a groupoid action $\beta : \mathbb{G} \rightarrow B(K \otimes H_G)$, where

$$H_G \stackrel{\text{def}}{=} \bigoplus_{w \in FP_r(\widehat{G})} (\mathbb{C} \cdot \xi_w)$$

is the Hilbert space with its Hilbert basis $\{\xi_w : w \in FP_r(\widehat{G})\}$, where

$$FP_r(\widehat{G}) \stackrel{\text{def}}{=} \mathbb{G} \setminus (V(\widehat{G}) \cup \{\emptyset\}).$$

We will call H_G the graph Hilbert space. A groupoid action β , called the graph-representation, is a nonunital intertwined partial representation, determining the bounded operators β_w on $K \otimes H_G$ satisfying that

$$\beta_w(m) R_w R_w^* = R_w^* m R_w,$$

and

$$\beta_v(m) = m,$$

for all $m \in M$, $w \in FP_r(\widehat{G})$ and $v \in V(\widehat{G})$, where R_w are the *right* multiplication operators on H_G with their symbols ξ_w , for all $w \in \mathbb{G}$. The adjoint R_w^* of R_w are defined to be $R_{w^{-1}}$, for all $w \in \mathbb{G}$.

Definition 2.1. *The crossed product algebra $\mathbb{M}_G = M \times_\beta \mathbb{G}$ is the von Neumann algebra generated by M and $\{R_w : w \in \mathbb{G}\}$, satisfying the above β -relations. We call $\mathbb{M}_G$ a right graph von Neumann algebra induced by G over M , via β . A right graph von Neumann algebra $\mathbb{M}_G$ has its canonical W^* -subalgebra $\mathbb{D}_G \stackrel{\text{def}}{=} \bigoplus_{v \in V(\widehat{G})} (M \cdot R_v)$, called the M -diagonal subalgebra of $\mathbb{M}_G$.*

Remark 2.1. *Recall that, in [10] and [11], we observed the (left) multiplication operators L_w 's, for all $w \in \mathbb{G}$, instead of using right multiplication operators R_w 's.*

Similar to [10], we construct an amalgamated W^* -probability space $(\mathbb{M}_G, E)$ over $\mathbb{D}_G$, where $E : \mathbb{M}_G \rightarrow \mathbb{D}_G$ is the canonical conditional expectation. Under this

setting, we realized that $\mathbb{M}_G$ is $*$ -isomorphic to a $\mathbb{D}_G$ -valued reduced free product $*_{\mathbb{D}_G}^r \mathbb{M}_e$ of the $\mathbb{D}_G$ -free blocks $\mathbb{M}_e$ indexed by directed edges $e \in E(G)$, where $e \in E(G)$

$$\mathbb{M}_e \stackrel{def}{=} vN(M \times_{\beta} \mathbb{G}_e, \mathbb{D}_G), \text{ for all } e \in E(G).$$

Here, $\mathbb{G}_e$ is a substructure (or a subgroupoid) of $\mathbb{G}$, consisting of all reduced words in $\{e, e^{-1}\}$, with the inherited admissibility on $\mathbb{G}$, for all $e \in E(G)$. Notice that the reduction of the free product “ $*_{\mathbb{D}_G}^r$ ” is completely dependent upon the admissibility on the graph groupoid $\mathbb{G}$.

Also, similar to [11], we can characterize the $\mathbb{D}_G$ -free blocks $\mathbb{M}_e$ ’s of our right graph von Neumann algebra $\mathbb{M}_G$. Because of the setting, in fact, the results are almost same. Especially, we show that each $\mathbb{D}_G$ -free block $\mathbb{M}_e$ of $\mathbb{M}_G$ is $*$ -isomorphic to a certain von Neumann algebra contained in either $\mathcal{M}_{lp}^{\beta}$ or $\mathcal{M}_{non-lp}^{\beta}$, where

$$\mathcal{M}_{lp}^{\beta} = \{vN(M \times_{\lambda_e} \mathbb{Z}, \mathbb{D}_G) : \lambda_e = \beta|_{\mathbb{S}_e}, e \text{ is a loop edge}\}$$

and

$$\mathcal{M}_{non-lp}^{\beta} = \{vN(M_2^{\beta_e}(M), \mathbb{D}_G) : e \text{ is a non-loop edge}\},$$

where $M \times_{\lambda(e)} \mathbb{Z}$ is a classical group crossed products of M and the infinite cyclic abelian group $\mathbb{Z}$ via a group action $\lambda^{(e)}$ satisfying that $\lambda_{(e)} = \beta|_{\mathbb{G}_e}$, and where $M_2^{\beta_e}(M)$ is a W^* -subalgebra of $M_2(M) = M \otimes_{\mathbb{C}} M_2(\mathbb{C})$ satisfying the β_e -relation: $\beta_e(m) R_e^* R_e^* m R_e = R_e^* m R_e$, for all $m \in M$ and for $e \in E(G)$, where $M_2(\mathbb{C})$ is the matricial algebra generated by all (2×2) -matrices. In particular, if $M = \mathbb{C}$, we can conclude that each $\mathbb{D}_G$ -free block $\mathbb{M}_e$ of $\mathbb{M}_G$ is $*$ -isomorphic to either $L^{\infty}(\mathbb{T})$ or $M_2(\mathbb{C})$, where $\mathbb{T}$ is the unit circle in $\mathbb{C}$, for all $e \in E(G)$. This characterization says that the study of right graph von Neumann algebras is the investigation of graph groupoids and the above two types of von Neumann algebras.

Assumption In this paper, we only consider the case where the fixed von Neumann algebra M is $\mathbb{C}$. Then every right graph von Neumann algebras $\mathbb{M}_G = \mathbb{C} \times_{\beta} \mathbb{G}$ are all $*$ -isomorphic to $M_G = \overline{\mathbb{C}[\mathbb{G}]}^w$, as W^* -subalgebras of $B(H_G)$, for any choice of β , by the linearity of β on $\mathbb{C}$. We will say that the von Neumann algebra M_G is the right graph von Neumann algebra of G . $\square$

We will consider the labeling operator τ of $\mathbb{G}$, which is an analogue of classical Heck-type operators (of groups). As an element of the right graph von Neumann algebra M_G , we can verify that τ has its decomposition $\tau = \oplus_{j=1}^N \tau_j$, where

$$N \stackrel{def}{=} \max\{\deg_{out}(v) : v \in V(\widehat{G})\}$$

and

$$\tau_j(\xi_w) \stackrel{def}{=} \begin{cases} \xi_w \xi_e = \xi_{we} & \text{if } we \neq \emptyset \text{ and} \\ & e \text{ has its weight } j \\ \xi_w \xi_\emptyset = \xi_\emptyset = 0 & \text{if there is no edge } e \text{ such that} \\ & e \text{ has its weight } j \text{ and } we \neq \emptyset. \end{cases}$$

The study of the labeling operator τ , itself, is interesting in Operator Theory and Quantum Physics, but we will concentrate on observing its free distributional data of the operator τ . The data will show how the graph groupoid $\mathbb{G}$ acts inside $B(H_G)$. Futhermore, if we consider the general case where we have $B(K \otimes H_G)$, for any Hilbert spaces K , we could show how $\mathbb{G}$ can acts on $K \otimes H_G$, too.

3. DEFINITIONS

In this chapter, we will introduce the definitions and concepts which we will use in the rest of this paper. We will review groupoids, groupoid actions, automata. Also, we observe the basic properties of right graph von Neumann algebras.

3.1. Categorical Groupoids and Groupoid Actions.

We say an algebraic structure $(\mathcal{X}, \mathcal{Y}, s, r)$ is a (*categorical*) *groupoid* if it satisfies that (i) $\mathcal{Y} \subset \mathcal{X}$, (ii) for all $x_1, x_2 \in \mathcal{X}$, there exists a partially-defined binary operation $(x_1, x_2) \mapsto x_1 x_2$, for all $x_1, x_2 \in \mathcal{X}$, depending on the source map s and the range map r satisfying that:

(ii-1) $x_1 x_2$ is well-determined, whenever $r(x_1) = s(x_2)$ and in this case, $s(x_1 x_2) = s(x_1)$ and $r(x_1 x_2) = r(x_2)$, for $x_1, x_2 \in \mathcal{X}$,

(ii-2) $(x_1 x_2) x_3 = x_1 (x_2 x_3)$, if they are well-determined in the sense of (ii-1), for $x_1, x_2, x_3 \in \mathcal{X}$,

(ii-3) if $x \in \mathcal{X}$, then there exist $y, y' \in \mathcal{Y}$ such that $s(x) = y$ and $r(x) = y'$, satisfying $x = y x y'$ (Here, the elements y and y' are not necessarily distinct),

(ii-4) if $x \in \mathcal{X}$, then there exists a unique element x^{-1} for x satisfying $x x^{-1} = s(x)$ and $x^{-1} x = r(x)$.

Thus, every group is a groupoid $(\mathcal{X}, \mathcal{Y}, s, r)$ with $|\mathcal{Y}| = 1$ (and hence $s = r$ on $\mathcal{X}$). This subset $\mathcal{Y}$ of $\mathcal{X}$ is said to be the *base of* $\mathcal{X}$. Remark that we can naturally assume that there exists the *empty element* $\emptyset$ in a groupoid $\mathcal{X}$. The empty element $\emptyset$ means the products $x_1 x_2$ are not well-defined, for some $x_1, x_2 \in \mathcal{X}$. Notice that if $|\mathcal{Y}| = 1$ (equivalently, if $\mathcal{X}$ is a group), then the empty word $\emptyset$ is not contained

in the groupoid $\mathcal{X}$. However, in general, whenever $|\mathcal{Y}| \geq 2$, a groupoid $\mathcal{X}$ always contain the empty word. So, if there is no confusion, we will naturally assume that the empty element $\emptyset$ is contained in $\mathcal{X}$.

It is easily checked that our graph groupoid $\mathbb{G}$ of a countable directed graph G is indeed a groupoid with its base $V(\widehat{G})$. i.e., every graph groupoid $\mathbb{G}$ of a countable directed graph G is a groupoid $(\mathbb{G}, V(\widehat{G}), s, r)$, where $s(w) = s(v \ w) = v$ and $r(w) = r(w \ v') = v'$, for all $w = v \ w \ v' \in \mathbb{G}$ with $v, v' \in V(\widehat{G})$. i.e., the vertex set $V(\widehat{G}) = V(G)$ is a base of $\mathbb{G}$.

Let $\mathcal{X}_k = (\mathcal{X}_k, \mathcal{Y}_k, s_k, r_k)$ be groupoids, for $k = 1, 2$. We say that a map $f : \mathcal{X}_1 \rightarrow \mathcal{X}_2$ is a *groupoid morphism* if (i) f is a function, (ii) $f(\mathcal{Y}_1) \subseteq \mathcal{Y}_2$, (iii) $s_2(f(x)) = f(s_1(x))$ in $\mathcal{X}_2$, for all $x \in \mathcal{X}_1$, and (iv) $r_2(f(x)) = f(r_1(x))$ in $\mathcal{X}_2$, for all $x \in \mathcal{X}_1$. If a groupoid morphism f is bijective, then we say that f is a *groupoid-isomorphism*, and the groupoids $\mathcal{X}_1$ and $\mathcal{X}_2$ are said to be *groupoid-isomorphic*.

Notice that, if two countable directed graphs G_1 and G_2 are graph-isomorphic, via a graph-isomorphism $g : G_1 \rightarrow G_2$, in the sense that (i) g is bijective from $V(G_1)$ onto $V(G_2)$, (ii) g is bijective from $E(G_1)$ onto $E(G_2)$, (iii) $g(e) = g(v_1 \ e \ v_2) = g(v_1) \ g(e) \ g(v_2)$ in $E(G_2)$, for all $e = v_1 \ e \ v_2 \in E(G_1)$, with $v_1, v_2 \in V(G_1)$, then the graph groupoids $\mathbb{G}_1$ and $\mathbb{G}_2$ are groupoid-isomorphic. More generally, if two graphs G_1 and G_2 have graph-isomorphic shadowed graphs $\widehat{G}_1$ and $\widehat{G}_2$, then $\mathbb{G}_1$ and $\mathbb{G}_2$ are groupoid-isomorphic.

Let $\mathcal{X} = (\mathcal{X}, \mathcal{Y}, s, r)$ be a groupoid. We say that this groupoid $\mathcal{X}$ *acts on a set* Y if there exists a groupoid action π of $\mathcal{X}$ such that $\pi(x) : Y \rightarrow Y$ is a well-determined function, for all $x \in \mathcal{X}$. Sometimes, we call the set Y , a $\mathcal{X}$ -*set*. We are interested in the case where a $\mathcal{X}$ -set Y is a Hilbert space. The nicest example of a groupoid action acting on a Hilbert space is a graph-representation in the sense of [10] and [11]. (Also, see [14] and [15] and the action β in Section 2.5.)

Let $\mathcal{X}_1 \subset \mathcal{X}_2$ be a subset, where $\mathcal{X}_2 = (\mathcal{X}_2, \mathcal{Y}_2, s, r)$ is a groupoid, and assume that $\mathcal{X}_1 = (\mathcal{X}_1, \mathcal{Y}_1, s, r)$, itself, is a groupoid, where $\mathcal{Y}_1 = \mathcal{X}_2 \cap \mathcal{Y}_2$. Then we say that the groupoid $\mathcal{X}_1$ is a *subgroupoid* of $\mathcal{X}_2$.

Recall that we say that a countable directed graph G_1 is a *full-subgraph* of a countable directed graph G_2 , if

$$E(G_1) \subseteq E(G_2)$$

and

$$V(G_1) = \{v \in V(G_1) : e = v \ e \text{ or } e = e \ v, \forall e \in E(G_1)\}.$$

Remark the difference between full-subgraphs and subgraphs: We say that G'_1 is a *subgraph* of G_2 , if

$$V(G'_1) \subseteq V(G_2)$$

and

$$E(G'_1) = \{e \in E(G_2) : e = v_1 e v_2, \text{ for } v_1, v_2 \in V(G'_1)\}.$$

We can see that the graph groupoid $\mathbb{G}_1$ of G_1 is a subgroupoid of the graph groupoid $\mathbb{G}_2$ of G_2 , whenever G_1 is a full-subgraph of G_2 .

3.2. Automata and Fractal Groups.

Automata Theory is the study of abstract machines, and we are using it in the formulation given by von Neumann. It is related to the theory of formal languages. In fact, automata may be thought of as the class of formal languages they are able to recognize. In von Neumann's version, an automaton is a finite state machine (FSM). i.e., a machine with input of symbols, transitions through a series of states according to a transition function (often expressed as a table). The transition function tells the automata which state to go to next, given a current state and a current symbol. The input is read sequentially, symbol by symbol, for example as a tape with a word written on it, registered by the head of the automaton; the head moves forward over the tape one symbol at a time. Once the input is depleted, the automaton stops. Depending on the state in which the automaton stops, it is said that the automaton either accepts or rejects the input. The set of all the words accepted by the automaton is called the language of the automaton. For the benefit for the readers, we offer the following references for the relevant part of Automata Theory: [1], [35], [36], [37], [50] and [51].

Let the quadruple $\mathcal{A} = \langle D, Q, \varphi, \psi \rangle$ be given, where D and Q are sets and

$$\varphi : D \times Q \rightarrow Q \quad \text{and} \quad \psi : D \times Q \rightarrow D$$

are maps. We say that D and Q are the (finite) *alphabet* and the *state set* of $\mathcal{A}$, respectively and we say that φ and ψ are the *output function* and the *state transition function*, respectively. In this case, the quadruple $\mathcal{A}$ is called an *automaton*. If the map $\psi(\bullet, q)$ is bijective on D , for any fixed $q \in Q$, then we say that the automaton $\mathcal{A}$ is *invertible*. Similarly, if the map $\varphi(x, \bullet)$ is bijective on Q , for any fixed $x \in D$, then we say that the automaton $\mathcal{A}$ is *reversible*. If the automaton $\mathcal{A}$ is both invertible and reversible, then $\mathcal{A}$ is said to be *bi-reversible*.

To help visualize the use of automata, a few concrete examples may help. With some oversimplification, they may be drawn from the analysis and synthesis of input / output models in Engineering, often referred to as black box diagram: excitation variables, response variables, and intermediate variables (e.g., see [54] and [55]).

In our presentation above, the D (the chosen finite alphabet) often takes different forms on the side of input D_i and output D_o . In popular automata that models stimuli of organisms, the three sets input D_i , output D_o , and the state set Q , could be as in the following prototypical three examples:

Example 3.1. *Models stimuli of organisms:*

$$\begin{aligned}
 D_i &= \{\text{positive stimulus, negative stimulus}\}, \\
 D_o &= \{\text{reaction, no reaction}\}, \\
 \text{and} \\
 Q &= \left\{ \begin{array}{l} \text{reaction to last positive stimulus,} \\ \text{no reaction to last positive stimulus} \end{array} \right\}.
 \end{aligned}$$

Example 3.2. In a control model for say a steering mechanism in a vehicle:

$$\begin{aligned}
 D_i &= \{\text{right, left}\}, \\
 D_o &= \{\text{switch on, switch off}\} \text{ or } \{\text{lamp on, lamp off}\} \\
 \text{and} \\
 Q &= \left\{ \begin{array}{l} \text{right-turning direction signal} \\ \text{left-turning direction signal} \end{array} \right\}.
 \end{aligned}$$

Example 3.3. In a model for quantization in Signal Processing:


$$D = D_i = D_o = \left\{ \begin{array}{l} \text{assignments from a bit alphabet,} \\ \text{with the bits referring to the value} \\ \text{of pulses-in and pulses-out in a} \\ \text{signal processing algorithm} \end{array} \right\},$$

for example, on a discrete multiresolution (e.g., [50]), and

$$Q = \begin{array}{l} \text{a subset of the Cartesian product of} \\ \text{copies of } D, \text{ fixing finite number of} \\ \text{times, i.e., } D \times \dots \times D \end{array}$$

Recently, various algebraists have studied automata and the corresponding automata groups (Also, see [1], [22], [35] and [37]). We will consider a certain special case, where Q is a free semigroupoid of a shadowed graph.

Roughly speaking, a *undirected tree* is a connected simplicial graph without loop finite paths. Recall that a (undirected) graph is *simplicial*, if the graph has neither loop-edges nor multi-edges. A *directed tree* is a connected simplicial graph without loop finite paths, with directed edges. In particular, we say that a directed tree $\mathcal{T}_n$ is a *n-regular tree*, if $\mathcal{T}_n$ is rooted and one-flowed infinite directed tree, having the same out-degrees n , for all vertices. For example, the 2-regular tree $\mathcal{T}_2$ can be depicted by



Let $\mathcal{A} = \langle D, Q, \varphi, \psi \rangle$ be an automaton with $|D| = n$. Then, we can construct *automata actions* of $\mathcal{A}$ on $\mathcal{T}_n$. Let's fix $q \in Q$. Then the action of $\mathcal{A}_q$ is defined on the finite words D_* of D by

$$\mathcal{A}_q(x) \stackrel{def}{=} \varphi(x, q), \text{ for all } x \in D,$$

and recursively,

$$\mathcal{A}_q((x_1, x_2, \dots, x_m)) = \varphi(x_1, \mathcal{A}_q(x_2, \dots, x_m)),$$

for all $(x_1, \dots, x_m) \in D_*$, where

$$D_* \stackrel{def}{=} \bigcup_{m=1}^{\infty} \left(\left\{ (x_1, \dots, x_m) \in D^m \mid \begin{array}{l} x_k \in D, \text{ for all } \\ k = 1, \dots, m \end{array} \right\} \right).$$

Then the automata actions $\mathcal{A}_q$'s are acting on the n -regular tree $\mathcal{T}_n$. In other words, all images of automata actions are regarded as an elements in the free semigroupoid $\mathbb{F}^+(\mathcal{T}_n)$ of the n -regular tree. i.e.,

$$V(\mathcal{T}_n) \supseteq D_*$$

and its edge set


$$E(\mathcal{T}_n) \supseteq \{\mathcal{A}_q(x) : x \in D, q \in Q\}.$$

This makes us to illustrate how the automata actions work.

Let $\mathcal{C} = \{\mathcal{A}_q : q \in Q\}$ be the collection of automata actions of the given automaton $\mathcal{A} = \langle D, Q, \varphi, \psi \rangle$. Then we can create a group $G(\mathcal{A})$ generated by the collection $\mathcal{C}$. This group $G(\mathcal{A})$ is called the *automata group generated by $\mathcal{A}$* . The generator set $\mathcal{C}$ of $G(\mathcal{A})$ acts “fully” on the $|D|$ -regular tree $\mathcal{T}_{|D|}$, we say that this group $G(\mathcal{A})$ is a *fractal group*. There are many ways to define fractal groups, but we define them in the sense of automata groups. (See [1] and [37]. In fact, in [37],

Batholdi, Grigorchuk and Nekrashevych did not define the term “fractal”, but they provide the fractal properties.)

Now, we will define a fractal group more precisely (Also see [1]). Let $\mathcal{A}$ be an automaton and let $\Gamma = G(\mathcal{A})$ be the automata group generated by the automata actions acting on the n -regular tree $\mathcal{T}_n$, where n is the cardinality of the alphabet of $\mathcal{A}$. By $St_\Gamma(k)$, denote the subgroup of $\Gamma = G(\mathcal{A})$, consisting of those elements of Γ , acting trivially on the k -th level of $\mathcal{T}_n$, for all $k \in \mathbb{N} \cup \{0\}$.




Analogously, for a vertex u in $\mathcal{T}_n$, define $St_\Gamma(u)$ by the subgroup of Γ , consisting of those elements of Γ , acting trivially on u . Then

$$St_\Gamma(k) = \bigcap_{u : \text{vertices of the } k\text{-th level of } \mathcal{T}_n} (St_\Gamma(u)).$$

For any vertex u of $\mathcal{T}_n$, we can define the algebraic projection $p_u : St_\Gamma(u) \rightarrow \Gamma$.


Definition 3.1. Let $\Gamma = G(\mathcal{A})$ be the automata group given as above. We say that this group Γ is a fractal group if, for any vertex u of $\mathcal{T}_n$, the image of the projection $p_u(St_\Gamma(u))$ is group-isomorphic to Γ , after the identification of the tree $\mathcal{T}_n$ with its subtree $\mathcal{T}_u$ with the root u .

For instance, if u is a vertex of the 2-regular tree $\mathcal{T}_2$, then we can construct a subtree $\mathcal{T}_u$, as follows:



As we can check, the graphs $\mathcal{T}_2$ and $\mathcal{T}_u$ are graph-isomorphic. The above definition shows that if the automata actions $\mathcal{A}_q$'s of $\mathcal{A}$ are acting *fully* on $\mathcal{T}_n$, then the automata group $G(\mathcal{A})$ is a fractal group. There are lots of famous fractal groups, but we introduce the following example, for our purpose.

Example 3.4. Let $\mathcal{A} = \langle X_{2n}, F_n, \varphi, \psi \rangle$ be an automaton, where F_n is the free group with its generator set $X_{2n} = \{g_1^{\pm 1}, \dots, g_n^{\pm 1}\}$. Then the automata group $G(\mathcal{A})$ is group-isomorphic to F_n . It is easy to check that all elements in F_n acts fully on the $2n$ -regular tree $\mathcal{T}_{2n}$, and hence $G(\mathcal{A})$ is a fractal group. For example, if $n = 2$, then we can get the following 0-th and 1-st levels of $\mathcal{T}_4$:



where e_{F_2} is the group-identity of $F_2 = \langle g_1, g_2 \rangle$.

3.3. Right Graph Von Neumann algebras.

Let G be a countable directed graph with its graph groupoid $\mathbb{G}$. We define and consider certain operators on the graph Hilbert space H_G induced by G . As we defined and observed, we can regard all elements in $\mathbb{G}$ as reduced words in $E(\widehat{G})$, under the admissibility with the reduction (RR).

Definition 3.2. Let G be a countable directed graph and let $\mathbb{G}$ be the corresponding graph groupoid. Define the Hilbert space H_G of G by

$$H_G \stackrel{\text{def}}{=} \bigoplus_{w \in FP_r(\widehat{G})} (\mathbb{C}\xi_w)$$

with its Hilbert basis $\mathcal{B}_{H_G} = \{\xi_w : w \in FP_r(\widehat{G})\}$ in H_G , where $FP_r(\widehat{G})$ is the reduced finite path set.

We have the following multiplication rule on H_G :

$$\xi_{w_1}\xi_{w_2} = \begin{cases} \xi_{w_1w_2} & \text{if } w_1 w_2 \neq \emptyset \\ \xi_\emptyset = 0_{H_G} & \text{otherwise,} \end{cases}$$

for all $\xi_{w_1}, \xi_{w_2} \in \mathcal{B}_{H_G}$. Suppose $w_1 = w$ and $w_2 = w^{-1}$ in $FP_r(\widehat{G})$. Then, by the above multiplication rule, we can have an element $\xi_w \xi_{w^{-1}} = \xi_{ww^{-1}}$ in H_G , where $w w^{-1}$ is a vertex in $V(\widehat{G})$. So, we can determine $\xi_v \in H_G$, for all $v \in V(\widehat{G})$, too. Thus, for any $w \in \mathbb{G}$, we can have the corresponding Hilbert space element ξ_w in H_G . This multiplication rule let us define left and right multiplication operators on H_G . The (left) multiplication operators L_w 's on H_G are defined by

$$L_w \xi_{w'} \stackrel{\text{def}}{=} \xi_w \xi_{w'} = \xi_{ww'}, \text{ for all } w, w' \in \mathbb{G},$$

where $\xi_\emptyset \stackrel{\text{def}}{=} 0_{H_G}$. It is easy to see that $L_{w_1} L_{w_2} = L_{w_1w_2}$, for all $w_1, w_2 \in \mathbb{G}$, and $L_w^* = L_{w^{-1}}$, for all $w \in \mathbb{G}$, and hence if w is a reduced finite path, then L_w is a partial isometry and if w is a vertex, then L_w is a projection on H_G . As in [13], we are interested in right multiplication operators:

Definition 3.3. An operator R_w on H_G is defined by the right multiplication operator with its symbol ξ_w on H_G , for $w \in \mathbb{G}$. i.e.,

$$R_w \xi_{w'} \stackrel{\text{def}}{=} \begin{cases} \xi_{w'} \xi_w = \xi_{w'w} & \text{if } w' w \neq \emptyset \\ \xi_\emptyset = 0_{H_G} & \text{otherwise,} \end{cases}$$

for all $w, w' \in \mathbb{G}$. The adjoint R_w^* of R_w is defined by $R_w^* = R_{w^{-1}}$, for all $w \in \mathbb{G}$.

By definition, the product of two right multiplication operators $R_{w_1} R_{w_2}$ is the multiplication operator $R_{w_2w_1}$. i.e.,

$$R_{w_1} R_{w_2} = R_{w_2w_1}, \text{ for all } w_1, w_2 \in \mathbb{G}.$$

So, it is easy to check that the multiplication operators R_v , for all $v \in V(\widehat{G})$, are projections on H_G , since

$$R_v^2 = R_v R_v = R_{v^2} = R_v \text{ and } R_v^* = R_{v^{-1}} = R_v.$$

Thus the right multiplication operators R_w , for all $w \in FP_r(\widehat{G})$, are partial isometries; indeed,

$$R_w R_w^* R_w = R_{ww^{-1}w} = R_w$$

and

$$R_w^* R_w R_w^* = R_{w^{-1}ww^{-1}} = R_{w^{-1}} = R_w^*,$$

for all $w \in FP_r(\widehat{G})$.

Now, we define certain groupoid actions β of the graph groupoid $\mathbb{G}$ on $K \otimes H_G$, called *right graph-representations*, where K is a Hilbert space.

Definition 3.4. Let $\mathbb{G}$ be a graph groupoid of a countable directed graph G and let M be a von Neumann algebra acting on a Hilbert space K (i.e., $M \subseteq B(K)$). Define a right graph-representation (in short, a right G -representation) $\beta : \mathbb{G} \rightarrow B(K \otimes H_G)$, by a nonunital intertwined partial representation satisfying that

$$\beta_w(m) R_w R_w^* = R_w^* m R_w = R_{w^{-1}} m R_w,$$

for all $m \in M$ and $w \in \mathbb{G}$, and

$$\beta_v(m) = m, \text{ for all } m \in M \text{ and } v \in V(\widehat{G}).$$

In this setting, we can regard R_w 's as $1_K \otimes R_w$'s on $K \otimes H_G$, for all $w \in \mathbb{G}$. The above two rules for β are called the β -relations on M .

For a fixed von Neumann algebra M and a graph groupoid $\mathbb{G}$, we can construct a groupoid crossed product algebra $M \times_\beta \mathbb{G}$.

Definition 3.5. Let M be a von Neumann algebra and $\mathbb{G}$, the graph groupoid of a countable directed graph G and let β be a right G -representation. Define the crossed product $\mathbb{M}_G = M \times_\beta \mathbb{G}$ of M and $\mathbb{G}$, via β , by the von Neumann algebra generated by M and $\{R_w : w \in \mathbb{G}\}$ in $B(K \otimes H_G)$, satisfying the β -relations on M . This von Neumann algebra $\mathbb{M}_G$ is called a right graph von Neumann algebra induced by G over M .

Remark 3.1. In [10] and [11], we defined a graph von Neumann algebra $\mathbb{M}_G^{(left)}$ by the groupoid crossed product $M \times_\alpha \mathbb{G}$, where $\alpha : \mathbb{G} \rightarrow B(K \otimes H_G)$ is a G -representation satisfying the α -relations on M :

$$\alpha_w(m) L_w L_w^* = L_w^* m L_w, \text{ for all } w \in FP_r(\widehat{G}),$$

and

$$\alpha_v(m) = m, \text{ for all } v \in V(\widehat{G}),$$

for all $m \in M$, where L_w 's are the (left) multiplication operators, for $w \in \mathbb{G}$. Notice that, in fact, the von Neumann algebra $\mathbb{M}_G^{(left)}$ is the opposite algebra $\mathbb{M}_G^{op}$ of a right graph von Neumann algebra $\mathbb{M}_G = M \times_\beta \mathbb{G}$, and hence they are anti- $*$ -isomorphic from each other. This guarantees that all results in [10] and [11] can be applicable in our “right” graph von Neumann algebraic setting.

Every operator x in a right graph von Neumann algebra $\mathbb{M}_G = M \times_\beta \mathbb{G}$ has its expression,

$$x = \sum_{w \in \mathbb{G}} m_w R_w, \text{ for } m_w \in M.$$

Lemma 3.1. *Let $m_1 R_{w_1}, \dots, m_n R_{w_n}$ be operators in a right graph von Neumann algebra $\mathbb{M}_G$, for $n \in \mathbb{N}$. Then*

$$\begin{aligned} & \Pi_{k=1}^n (m_k R_{w_k}) \\ &= \begin{cases} \left(m_1 m_2^{w_1^{-1}} m_3^{(w_1 w_2)^{-1}} \dots m_n^{(w_1 w_2 \dots w_{n-1})^{-1}} \right) R_{w_n \dots w_1} & \text{if } w_n \dots w_1 \neq \emptyset \\ 0_{\mathbb{M}_G} & \text{otherwise,} \end{cases} \end{aligned}$$

where $m^w \stackrel{\text{def}}{=} \beta_w(m)$, for all $m \in M$ and $w \in \mathbb{G}$. $\square$

3.4. M -Valued Right Graph W^* -Probability Spaces.

Let G be a countable directed graph with its graph groupoid $\mathbb{G}$ and let M be an arbitrary von Neumann algebra acting on a Hilbert space K . In this section, we will define a M -diagonal right graph W^* -probability space $(\mathbb{M}_G, E)$, over its M -diagonal subalgebra $\mathbb{D}_G$, where $\mathbb{M}_G = M \times_\beta \mathbb{G}$ is a right graph von Neumann algebra.

Let $v \in V(\widehat{G})$. Then we can define a conditional expectation $E_v : \mathbb{M}_G \rightarrow M \cdot R_v$, where $M \cdot R_v = \{m R_v : m \in M\}$, by

$$E_v \left(\sum_{w \in \mathbb{G}} m_w R_w \right) = m_v R_v,$$

for all $\sum_{w \in \mathbb{G}} m_w R_w \in \mathbb{M}_G$. Notice that each $M \cdot R_v$ is a W^* -subalgebra of $\mathbb{M}_G$, which is $*$ -isomorphic to M , since R_v is a projection, for all $v \in V(\widehat{G})$. A pair $(\mathbb{M}_G, E_v)$ is a M -valued W^* -probability space, for $v \in V(\widehat{G})$. We call it a *vertex-depending*

(or the v -depending) *right graph W^* -probability space over M* . The conditional expectation E_v is said to be a *vertex-depending* (or the v -depending) *conditional expectation for $v \in V(\widehat{G})$* . By the very definition, the right graph von Neumann algebra $\mathbb{M}_G$ has $|V(\widehat{G})|$ -many vertex-depending W^* -probability spaces over M .

Definition 3.6. By $\mathbb{D}_G$, denote the W^* -subalgebra $\bigoplus_{v \in V(\widehat{G})} (M \cdot R_v)$ of $\mathbb{M}_G = M \times_\beta \mathbb{G}$. This subalgebra $\mathbb{D}_G$ is called the *M -diagonal subalgebra of $\mathbb{M}_G$* . Define a conditional expectation $E : \mathbb{M}_G \rightarrow \mathbb{D}_G$, by $\bigoplus_{v \in V(\widehat{G})} E_v$, where E_v 's are the v -depending conditional expectations, for all $v \in V(\widehat{G})$. i.e.,

$$E \left(\sum_{w \in \mathbb{G}} m_w R_w \right) = \sum_{v \in V(\widehat{G})} m_v R_v,$$

for all $\sum_{w \in \mathbb{G}} m_w R_w \in \mathbb{M}_G$. The pair $(\mathbb{M}_G, E)$ is called the *M -diagonal right graph W^* -probability space over $\mathbb{D}_G$* . We call E , the *canonical conditional expectation*.

Notice that the M -diagonal subalgebra $\mathbb{D}_G$ of $\mathbb{M}_G = M \times_\beta \mathbb{G}$ is uniquely determined for all right G -representations β .

Definition 3.7. Let G be a countable directed graph and let $\mathbb{G}$ be the graph groupoid of G . Define a map $\delta : \mathbb{G} \rightarrow \mathbb{G}$ by mapping $w \in \mathbb{G}$ to the graphical image $\delta(w)$ of w , for all $w \in \mathbb{G}$. The map δ on $\mathbb{G}$ is called the *diagram map* and the graphical image $\delta(w)$ of w is called the *diagram of w* , for all $w \in \mathbb{G}$. We say that the elements w_1 and w_2 in $\mathbb{G}$ are *diagram-distinct*, if (i) $w_1 \neq w_2^{-1}$ and (ii) $\delta(w_1) \neq \delta(w_2)$. Suppose that X_1 and X_2 are subsets in $\mathbb{G}$. They are said to be *diagram-distinct* if, for any pair (w_1, w_2) in $X_1 \times X_2$, w_1 and w_2 are diagram-distinct.

Let l be a loop edge in $E(\widehat{G})$, and let $w_1 = l^{k_1}$ and $w_2 = l^{k_2}$ in $\mathbb{G}$, for $k_1 \neq k_2 \in \mathbb{N}$. Clearly, $w_1 \neq w_2^{-1}$. But the elements w_1 and w_2 are not diagram-distinct, because their diagrams $\delta(w_1)$ and $\delta(w_2)$ are identical to the diagram $\delta(l) = l$ of l . Suppose either w_1 or w_2 is a non-loop finite path such that (i) $w_1 \neq w_2^{-1}$, and (ii) $\delta(w_k)$'s are not loop finite paths, for $k = 1, 2$. Then they are diagram-distinct whenever w_1 and w_2 are distinct in $\mathbb{G}$.

Theorem 3.2. The subsets $M \cdot R_{w_1}$ and $M \cdot R_{w_2}$ of a right graph von Neumann algebra $\mathbb{M}_G = M \times_\beta \mathbb{G}$ are free over $\mathbb{D}_G$ in $(\mathbb{M}_G, E)$ if and only if w_1 and w_2 are diagram-distinct in $\mathbb{G}$, where $M \cdot R_w \stackrel{\text{def}}{=} \{m L_w : m \in M\}$, for all $w \in \mathbb{G}$. $\square$

The readers can find the proof showing that: $M \cdot L_{w_1}$ and $M \cdot L_{w_2}$ are free over $\mathbb{D}_G$ in a (left) graph von Neumann algebra $M \times_\alpha \mathbb{G}$, if and only if w_1 and w_2 are diagram-distinct, in [10] and [11], where L_{w_1} and L_{w_2} are left multiplication

operators. By the slight modification of it, we can prove the above theorem. As corollary of the previous theorem, if e_1 and e_2 are distinct edges in $E(G)$ ($\subset E(\widehat{G})$), then $M \cdot R_{e_1}$ and $M \cdot R_{e_2}$ are free over $\mathbb{D}_G$ in $(\mathbb{M}_G, E)$, and vice versa. Indeed, if two edges e_1 and e_2 are distinct in $E(G)$, then e_1 and $e_2^{\pm 1}$ (resp., e_2 and $e_1^{\pm 1}$) are diagram-distinct in $\mathbb{G}$.

Let's consider the subset $\{e, e^{-1}\}$ of $\mathbb{G}$, for $e \in E(G) \subset E(\widehat{G})$. Then we can define a subset $\mathbb{G}_e$ of $\mathbb{G}$, by the collection of all reduced words in $\{e, e^{-1}\}$. It is easy to check that: if e is a loop edge in $E(G)$, then $\mathbb{G}_e$ is a group, which is group-isomorphic to the infinite abelian cyclic group $\mathbb{Z}$, and if e is a non-loop edge in $E(G)$, then $\mathbb{G}_e$ is a groupoid $\{\emptyset, v_1, v_2, e, e^{-1}\}$, where $e = v_1 e v_2$, with $v_1, v_2 \in V(G)$.

Notice that the graph groupoid $\mathbb{G}$ is the reduced free product $\bigast_{e \in E(G)} \mathbb{G}_e$ of $\mathbb{G}_e$'s (in the sense of [10]). Moreover, each $\mathbb{G}_e$ can be regarded as a new graph groupoid induced by a one-edge graph G_e , with $E(G_e) = \{e\}$. Clearly, here, G_e is a full-subgraph of G , and $\mathbb{G}_e$ is a subgroupoid of $\mathbb{G}$, for all $e \in E(G)$. Construct the W^* -subalgebras,

$$\mathbb{M}_e \stackrel{\text{def}}{=} vN(M \times_{\beta_e} \mathbb{G}_e, \mathbb{D}_G) \text{ with } \beta_e \stackrel{\text{def}}{=} \beta|_{\mathbb{G}_e} \text{ on } M,$$

of $\mathbb{M}_G$, for all $e \in E(G)$. By the previous theorem, we can have that:

Corollary 3.3. *The W^* -subalgebras $\mathbb{M}_e$'s of $\mathbb{M}_G$ are free over $\mathbb{D}_G$, from each other, for all $e \in E(G)$, in $(\mathbb{M}_G, E)$. $\square$*

Definition 3.8. *The W^* -subalgebras $\mathbb{M}_e$ of $\mathbb{M}_G$ are called $\mathbb{D}_G$ -free blocks of $\mathbb{M}_G$, for all $e \in E(G)$.*

Notice that $\mathbb{M}_e$ is (not only $*$ -isomorphic but also) identically same as $\mathbb{M}_{e^{-1}}$ in $\mathbb{M}_G$, for all $e \in E(G)$. So, we can get the following theorem characterizing the $\mathbb{D}_G$ -free structure of $(\mathbb{M}_G, E)$. The following theorem is the direct consequence of the previous corollary, and the fact that $\{\mathbb{M}_e : e \in E(G)\}$ generates $\mathbb{M}_G$.

Theorem 3.4. *Let $(\mathbb{M}_G, E)$ be the M -diagonal right graph W^* -probability space over its M -diagonal subalgebra $\mathbb{D}_G$. Then*

$$\mathbb{M}_G \stackrel{* \text{-isomorphic}}{=} \bigast_{e \in E(G)}^{*\mathbb{D}_G} \mathbb{M}_e,$$

where $\mathbb{M}_e$'s are the $\mathbb{D}_G$ -free blocks of $\mathbb{M}_G$, for all $e \in E(G)$. $\square$

Recall that if (A, E_B) is a B -valued W^* -probability space over B , and if $\{A_i\}_{i \in I}$ are W^* -subalgebras of A , over their common W^* -subalgebra B , then the B -free product algebra $\ast_B \prod_{i \in I} A_i$ can be expressed by

$$B \oplus \left(\bigoplus_{n=1}^{\infty} \left(\bigoplus_{i_1 \neq i_2, i_2 \neq i_3, \dots, i_{n-1} \neq i_n \in I} (A_{i_1}^o \otimes_B \dots \otimes_B A_{i_n}^o) \right) \right),$$

as a Banach space, where $A_{i_k}^o \stackrel{\text{def}}{=} A_{i_k} \ominus B$, for all $k = 1, \dots, n$. Since our right graph von Neumann algebra $\mathbb{M}_G$ is $\ast$ -isomorphic to the $\mathbb{D}_G$ -free product $\ast_{\mathbb{D}_G} \prod_{e \in E(G)} \mathbb{M}_e$, where $\mathbb{M}_e$'s are the $\mathbb{D}_G$ -free blocks of $\mathbb{M}_G$, we can get a similar Banach space expression. However, the $\mathbb{D}_G$ -tensor products in our $\mathbb{D}_G$ -free product algebra is dependent upon the admissibility on $\mathbb{G}$. i.e., if two edges e_1 and e_2 are totally distinct, in the sense that all pairs (e_1, e_2) are diagram-distinct, for $e_k \in \{e_k, e_k^{-1}\}$, for $k = 1, 2$, then $\mathbb{M}_{e_1}^o \otimes_{\mathbb{D}_G} \mathbb{M}_{e_2}^o$ is Banach-space isomorphic to $\{0_{\mathbb{D}_G}\}$, where $\mathbb{M}_{e_k}^o = \mathbb{M}_{e_k} \ominus \mathbb{D}_G$, for $k = 1, 2$.

Definition 3.9. Let G be a countable directed graph. Define the subset $E(\widehat{G})_r^*$ of $FP_r(\widehat{G})$ by

$$E(\widehat{G})_r^* \stackrel{\text{def}}{=} E(\widehat{G}) \cup \left(\bigcup_{k=2}^{\infty} \left\{ e_1 e_2 \dots e_{k-1} e_k \left| \begin{array}{l} e_1 \dots e_k \in FP_r(\widehat{G}), \\ e_1 \neq e_2^{\pm 1}, e_2 \neq e_3^{\pm 1}, \\ \dots, e_{k-1} \neq e_k^{\pm 1} \end{array} \right. \right\} \right).$$

Similar to [10] and [11], we can get the following theorem.

Theorem 3.5. Let G be a countable directed graph and let $\mathbb{M}_G = M \times_{\beta} \mathbb{G}$ be the corresponding right graph von Neumann algebra, for a fixed von Neumann algebra M . Then, as a Banach space,

$$\mathbb{M}_G = \mathbb{D}_G \oplus \left(\bigoplus_{w \in E(\widehat{G})_r^*} \mathbb{M}_w^o \right)$$

with

$$\mathbb{M}_w^o \stackrel{\text{def}}{=} \mathbb{M}_{e_1}^o \otimes_{\mathbb{D}_G} \dots \otimes_{\mathbb{D}_G} \mathbb{M}_{e_n}^o,$$

where

$$\mathbb{M}_e^o \stackrel{\text{def}}{=} \mathbb{M}_e \ominus \mathbb{D}_G, \text{ for all } e \in E(\widehat{G}),$$

whenever $w = e_1 \dots e_n \in E(\widehat{G})_r^*$, with $e_1, \dots, e_n \in E(\widehat{G})$, under the identification $\mathbb{M}_e = M_{e^{-1}}$, for all $e \in E(\widehat{G})$. $\square$

Notation To emphasize that the $\mathbb{D}_G$ -valued free product algebra $\ast_{\mathbb{D}_G} \prod_{e \in E(G)} \mathbb{M}_e$ has the above Banach-space expression determined by the admissibility on $\mathbb{G}$, we

will denote it by $\ast_{\mathbb{D}_G}^r \mathbb{M}_e$. And we will call it the $\mathbb{D}_G$ -valued *reduced* free product of the $\mathbb{D}_G$ -free blocks $\mathbb{M}_e$'s. $\square$

In [11], the $\mathbb{D}_G$ -free blocks in a (left) graph von Neumann algebra are characterized. Similarly, we can have the following results.

Theorem 3.6. *Let $\mathbb{M}_G = M \times_\beta \mathbb{G}$ be a right graph von Neumann algebra. Then its $\mathbb{D}_G$ -free block $\mathbb{M}_e$ is $\ast$ -isomorphic to either $M \times_{\lambda_e} \mathbb{Z}$ (whenever e is a loop-edge) or $M \otimes_{\beta_e} M_2(\mathbb{C})$ (whenever e is a non-loop edge), where λ_e is the group action of $\mathbb{Z}$, such that $\lambda_n(m) = \beta_{e^n}(m)$, for all $m \in M$ and $n \in \mathbb{Z}$. $\square$*

Assumption In the rest of this paper, we will restrict our interests to the case where $M = \mathbb{C}$. And we call a graph von Neumann algebra $\mathbb{M}_G = \mathbb{C} \times_\beta \mathbb{G}$, “the” right graph von Neumann algebra induced by G . $\square$

Suppose $M = \mathbb{C}$. Then a right graph von Neumann algebra $\mathbb{M}_G = \mathbb{C} \times_\beta \mathbb{G}$ is $\ast$ -isomorphic to the groupoid $W^\ast$ -subalgebra $\overline{\mathbb{C}[\mathbb{G}]}^w$ of $B(H_G)$, for all right graph-representation β , where H_G is the graph Hilbert space. Also, in such case, the $\mathbb{D}_G$ -free blocks $\mathbb{M}_e$'s of $\mathbb{M}_G$ are $\ast$ -isomorphic to either

$$L(\mathbb{Z}) \stackrel{\ast\text{-isomorphic}}{=} L^\infty(\mathbb{T}) \quad \text{or} \quad M_2(\mathbb{C}).$$

Example 3.5. *Let G_N be a one-vertex- N -loop-edge graph with its vertex set $V(G_N) = \{v\}$ and $E(G_N) = \{e_j = v e_j v : j = 1, \dots, N\}$. Then the right graph von Neumann algebra $\mathbb{C}_{G_N} = \mathbb{C} \times_\beta \mathbb{G}_N$, where $\mathbb{G}_N$ is the graph groupoid of G_N , is $\ast$ -isomorphic to the free group factor $L(F_N)$ (Also See [10]). Notice that the graph groupoid $\mathbb{G}_N$ of G_N is a group and moreover it is group-isomorphic to the free group $F_N = \langle g_1, \dots, g_N \rangle$ with N -generators $\{g_1, \dots, g_N\}$. Indeed, the edges $e_1, \dots, e_N$ are generators of $\mathbb{G}_N$ as a group. So, there is a natural generator-preserving group-isomorphism between $\mathbb{G}_N$ and F_N . Also, notice that the vertex v in $\mathbb{G}_N$ is the group identity. And all partial isometries $R_{e_j^n}$ correspond to the unitary representation $u_{g_j^n}$ of g_j^n , for all $n \in \mathbb{Z} \setminus \{0\}$ and for $j = 1, \dots, N$. Therefore, the right graph von Neumann algebra $\mathbb{C}_{G_N} = \mathbb{C} \times_\beta \mathbb{G}_N$ is $\ast$ -isomorphic to the classical crossed product algebra $\mathbb{C} \times_\lambda F_N$, whenever $\lambda = \beta$. Notice that $\mathbb{C} \times_\beta \mathbb{G}_N$ is $\ast$ -isomorphic to $\overline{\mathbb{C}[\mathbb{G}_N]}^w$, for any β , by the linearity of β on $\mathbb{C}$. Also, it is well-known that a crossed product algebra $\mathbb{C} \times_\gamma \Gamma$ is $\ast$ -isomorphic to $\overline{\mathbb{C}[\Gamma]}^w$, for any group action γ , for all groups Γ . Therefore, the right graph von Neumann algebra $\mathbb{C}_{G_N}$ is $\ast$ -isomorphic to the group von Neumann algebra $\overline{\mathbb{C}[F_N]}^w$. More generally, the right graph von Neumann algebra $\mathbb{M}_{G_N} = M \times_\beta \mathbb{G}_N$ is $\ast$ -isomorphic to the classical crossed product von Neumann algebra $M \times_\lambda F_N$, whenever $\lambda = \beta$.*

Example 3.6. *Let C_N be the one-flow circulant graph with $V(C_N) = \{v_1, \dots, v_N\}$ and $E(C_N) = \{e_j = v_j e_j v_{j+1} : j = 1, \dots, N, v_{N+1} \stackrel{\text{def}}{=} v_1\}$. Then the right graph von Neumann algebra $\mathbb{M}_{C_N} = \overline{\mathbb{C}[\mathbb{G}_{C_N}]}^w$ contains $W^\ast$ -subalgebras which are*

$\ast$ -isomorphic to the group von Neumann algebra $L(\mathbb{Z}) = \overline{\mathbb{C}[\mathbb{Z}]}^w$. Since a finite path $w = e_1 \dots e_N$ induces the subset $\mathbb{S}_w$ of $\mathbb{G}_{C_N}$ consisting of all reduced words in $S_w = \{w, w^{-1}\}$ in $\mathbb{G}_{C_N}$, we have the $W^\ast$ -subalgebra $N = \overline{\mathbb{C}(\mathbb{S}_w)}^w$ in $\mathbb{M}_{C_N}$. We can regard this $W^\ast$ -subalgebra N as the right graph von Neumann algebra induced by the graph G_w with its vertex set $V(G_w) = \{v_1\}$ and its edge set $E(G_w) = \{w = v_1 w v_1\}$. Then it is an one-vertex-one-loop-edge graph. So, by the previous example, the von Neumann algebra N is $\ast$ -isomorphic to $L(\mathbb{Z})$, where $\mathbb{Z}$ is the infinite abelian cyclic group. It is well-known that the group von Neumann algebra $L(\mathbb{Z})$ is $\ast$ -isomorphic to $L^\infty(\mathbb{T})$, where $\mathbb{T}$ is the unit circle in $\mathbb{C}$. Consider w^k , for all $k \in \mathbb{Z} \setminus \{0\}$. Then $W^\ast$ -subalgebras $N = \mathbb{C} \times_\beta \mathbb{G}_{w^k}$ of $\mathbb{C}_{C_N}$ are all $\ast$ -isomorphic to $L(\mathbb{Z})$.

We can easily check that $E(C_N)_r^\ast = FP(C_N) \cup FP(C_N^{-1})$, for all $N \in \mathbb{N}$. Therefore,

$$\mathbb{M}_{C_N} = \ast_{k=1}^N \mathbb{M}_{e_k}$$

where $\mathbb{M}_{e_k} \stackrel{\text{def}}{=} vN \left(\overline{\mathbb{C}(\mathbb{G}_e)}^w, \mathbb{D}_{C_N} \right)$, for all $k = 1, \dots, N$

$$\begin{aligned} &= \mathbb{D}_{C_N} \oplus \left(\bigoplus_{w^\ast \in E(C_N)_r^\ast} \mathbb{M}_{w^\ast}^o \right) \\ &= \mathbb{D}_{C_N} \oplus \left(\bigoplus_{w \in FP(C_N) \cup FP(C_N^{-1})} \mathbb{M}_w^o \right), \end{aligned}$$

where $\mathbb{M}_w^o \stackrel{\text{def}}{=} \mathbb{M}_{e^{(1)}}^o \otimes_{\mathbb{D}_{C_N}} \dots \otimes_{\mathbb{D}_{C_N}} \mathbb{M}_{e^{(n)}}^o$, whenever $w = e^{(1)} \dots e^{(n)}$ in $E(C_N)_r^\ast$, for $n \in \mathbb{N}$.

4. LABELED GRAPH GROUPOIDS AND GRAPH AUTOMATA

We will give suitable weights to the elements of graph groupoids induced by the given locally finite connected directed graphs. Then these weights give more accurate information of the admissibility on graph groupoids. Such admissibility conditions are explained by the labeling map and the shifting (map) of the automata constructed by graph groupoids. Also, this process would show how the admissibility on a graph groupoid works on the corresponding graph Hilbert space.

Recall that a countable directed graph G is *locally finite*, if every vertex of G has only finitely many incident edges (equivalently, the degree of each vertex is finite). Also, we say that G is *connected*, if, for any pair (v_1, v_2) of distinct vertices $(v_1 \neq$

v_2), there always exists at least one reduced finite path $w \in FP_r(\widehat{G})$ such that $w = v_1 w v_2$ and $w^{-1} = v_2 w^{-1} v_1$.

Assumption From now on, all given directed graphs are locally finite and connected. $\square$

Let G be a locally finite connected directed graph and let v_0 be an any fixed vertex of G . Then we can define the out-degree $\deg_{out}(v_0)$, the in-degree $\deg_{in}(v_0)$, and the degree $\deg(v_0)$ of v_0 as follows:

$$\deg_{out}(v_0) \stackrel{def}{=} |\{e \in E(G) : e = v_0 e\}|,$$

$$\deg_{in}(v_0) \stackrel{def}{=} |\{e \in E(G) : e = e v_0\}|,$$

and

$$\deg(v_0) \stackrel{def}{=} \deg_{out}(v_0) + \deg_{in}(v_0).$$

By the locally-finiteness of G , the above three cardinalities are less than ∞ . We are interested in out-degrees.

4.1. Labeled Graph Groupoids.

Let G be a locally finite connected directed graph. By the locally finiteness of G , we can have that

$$\max\{\deg_{out}(v) : v \in V(G)\} < \infty.$$

Denote the maximal value of the out-degrees of all vertices by N . Then we can define a set

$$Y = \{1, \dots, N\} \text{ and } Y_0 = \{0\} \cup Y.$$

Define *lattices* $l_0, l_1, \dots, l_N$ in the 2-dimensional real vector space $\mathbb{R}^2$, by the vectors

$$l_0 = \overrightarrow{(0, 0)} \text{ and } l_k = \overrightarrow{(1, e^k)},$$

for all $k = 1, \dots, N$. To distinguish the vectors and points in $\mathbb{R}^2$, we use the notation $\overrightarrow{(t_1, t_2)}$, as a vector connecting $(0, 0)$ and (t_1, t_2) in $\mathbb{R}^2$, and (t_1, t_2) , as a point in $\mathbb{R}^2$, for $t_1, t_2 \in \mathbb{R}$. For the given lattices $l_0, l_1, \dots, l_N$, determine the *inverse lattices* $l_{-0}, l_{-1}, \dots, l_{-N}$ by

$$l_{-0} = \overrightarrow{(0,0)} = l_0, \text{ and } l_{-k} = \overrightarrow{(1, -e^k)},$$

for all $k = 1, \dots, N$. Since $l_{-0} = l_0$, it is the self-inverse lattice. i.e., we can understand the lattices $l_1, \dots, l_N$ are the upward lattices in $\mathbb{R}^2$, and the inverse lattices $l_{-1}, \dots, l_{-N}$ are the downward lattices in $\mathbb{R}^2$.

Define now the *lattice paths on $\mathbb{R}^2$, generated by the lattices $l_{\pm 1}, \dots, l_{\pm N}$* , by the diagrams in $\mathbb{R}^2$, constructed by the rule:

$l_i l_j$ = the diagram connecting l_i and l_j by identifying the end point $(1, t_i e^i)$ of l_i and the starting point $(0, 0)$ of the l_j ,

where

$$t_i \stackrel{\text{def}}{=} \begin{cases} 1 & \text{if } i \in \{1, \dots, N\} \\ -1 & \text{if } i \in \{-1, \dots, -N\}, \end{cases}$$

for all $i \in \{\pm 1, \dots, \pm N\}$. For instance, if $l_1 = (0, e)$ and $l_{-3} = (0, -e^3)$, then a lattice path $l_1 l_{-3}$ is the diagram in $\mathbb{R}^2$, starting at $(0, 0)$ and ending at $(2, e - e^3)$, via $(1, e)$. We can do the above process inductively, and hence we can create the lattice paths $l_{i_1} l_{i_2} \dots l_{i_n}$ on $\mathbb{R}^2$, for all $(i_1, \dots, i_n) \in \{\pm 1, \dots, \pm N\}^n$, for $n \in \mathbb{N}$. Suppose we have the above lattice path $l_1 l_{-3}$, and let $l_2 = (1, e^2)$. Then, inductively, we can have the lattice path $l_1 l_{-3} l_2$, by identifying the ending point $(2, e - e^3)$ of $l_1 l_{-3}$ and the starting point $(0, 0)$ of l_2 . Then it is a lattice path starting at $(0, 0)$ and ending at $(3, e - e^3 + e^2)$, via $(1, e)$ and $(2, e - e^3)$ (Also, see [17] and [41]).

Let $l = l_{i_1} l_{i_2} \dots l_{i_n}$ be a lattice path generated by $l_{\pm 1}, \dots, l_{\pm N}$. Then the *length* $|l|$ of l is defined to be n , the cardinality of lattices generating l .

Definition 4.1. Let $N = \max\{\deg_{\text{out}}(v) : v \in V(G)\} \in \mathbb{N}$, and let $X = \{l_{\pm 1}, \dots, l_{\pm N}\}$ be the lattices defined as above. Define the set X_0 by $X \cup \{l_0\}$. The set $\mathcal{L}_N$ is the collection of all lattice paths generated by X , containing l_0 . Define a subset $\mathcal{L}_N(k)$ of $\mathcal{L}_N$ by

$$\mathcal{L}_N(k) \stackrel{\text{def}}{=} \{l \in \mathcal{L}_N : |l| = k\},$$

for all $k \in \mathbb{N}$.

By definition, it is clear that

$$\mathcal{L}_N = \{l_0\} \sqcup \left(\bigcup_{k=1}^{\infty} (\mathcal{L}_N(k)) \right),$$

where $\sqcup$ means the disjoint union. Suppose $l = l_{i_1} l_{i_2} \dots l_{i_n} \in \mathcal{L}_N(n)$, for $(i_1, \dots, i_n) \in \{\pm 1, \dots, \pm N\}^n$, for $n \in \mathbb{N}$. Also assume that this length- n lattice path l

starts at $(0, 0)$ and ends at $(0, 0)$ in $\mathbb{R}^2$. Then we say that the lattice path l satisfies the axis property. It is easy to check that:

Lemma 4.1. *If n is odd, then every length- n lattice path does not satisfy the axis property. $\square$*

Define now the subset $\mathcal{L}_N^o(k)$ of $\mathcal{L}_N(k)$ satisfying the axis property.

Definition 4.2. *Let $\mathcal{L}_N(k)$ be the set of all length- k lattice paths on $\mathbb{R}^2$. Define the subset $\mathcal{L}_N^o(k)$ of $\mathcal{L}_N(k)$ by*

$$\mathcal{L}_N^o(k) \stackrel{\text{def}}{=} \{l \in \mathcal{L}_N(k) : l \text{ satisfies the axis property}\},$$

for all $k \in \mathbb{N}$. And define the subset $\mathcal{L}_N^o$ of $\mathcal{L}_N$ by $\bigsqcup_{k=1}^{\infty} (\mathcal{L}_N^o(k))$.

Now, we label the elements of a graph groupoid $\mathbb{G}$ by X_0 (or $\mathcal{L}_N$). As above, let

$$X = \{l_1, \dots, l_N\} \text{ and } X_0 = \{l_0, l_1, \dots, l_N\},$$

where

$$l_0 = \overrightarrow{(0, 0)} \text{ and } l_k = \overrightarrow{(0, e^k)},$$

for all $j = 1, \dots, N = \max\{\deg_{out}(v) : v \in V(G)\}$.

Definition 4.3. *Let G be a given graph and let X and X_0 be the set of our lattices as above in $\mathbb{R}^2$. We call both X and $X_0 = \{l_0\} \cup X$, the (lattice) labeling sets of G .*

Let's fix $v_0 \in V(G)$, and let $\deg_{out}(v_0) = n \leq N$. Define the subset $E_{out}^{v_0}$ of the edge set $E(G)$ by

$$E_{out}^{v_0} \stackrel{\text{def}}{=} \{e \in E(G) : e = v_0 e\} \subseteq E(G).$$

i.e., $\deg_{out}(v_0) = |E_{out}^{v_0}|$. Then we decide the weights $\{(v_0, l_1), \dots, (v_0, l_n)\}$, contained in the set $\{v_0\} \times X$, on $E_{out}^{v_0}$. i.e.,

$$E_{out}^{v_0} = \left\{ e_{(v_0, l_1)}, \dots, e_{(v_0, l_n)} : \begin{array}{l} \omega_{v_0}(e_{(v_0, l_j)}) = (v_0, l_j), \\ \forall j = 1, \dots, n \end{array} \right\},$$

where

$$\omega_{v_0} : E_{out}^{v_0} \rightarrow \{v_0\} \times \{l_1, \dots, l_n\}$$

is the *weighting process*, putting the weight (v_0, l_j) to an edge $e_{(v_0, l_j)} \in E_{out}^{v_0}$.

For any $v \in V(G)$, we can do the same process and we have the corresponding weighting process ω_v and the corresponding set

$$E_{out}^v = \left\{ e_{(v,l_1)}, \dots, e_{(v,l_k)} \left| \begin{array}{l} l_1, \dots, l_k \in X \\ \text{and } \omega_v(e_{(v,l_j)}) = (v, l_j), \\ \forall j = 1, \dots, k, \text{ where} \\ k = \deg_{out}(v) \leq N \end{array} \right. \right\}.$$

Notice that if $\deg_{out}(v) = 0$, then E_{out}^v is empty, by the fact that $\{v\} \times \emptyset = \emptyset$, where $\emptyset$ is the empty set. Notice also that, by the connectedness of G ,

$$E(G) = \bigsqcup_{v \in V(G)} E_{out}^v,$$

where “ $\bigsqcup$ ” means the disjoint union. So, the each weighting processes $(\omega_v)_{v \in V(G)}$ extends to the *weighting process of $E(G)$* by X :

$$\omega : E(G) \rightarrow V(G) \times X$$

by

$$\omega(e) = \omega(v e) \stackrel{def}{=} \omega_v(e),$$

for all $e = v e \in E(G)$, with $v \in V(G)$. i.e., $\omega = \bigcup_{v \in V(G)} \omega_v$.

Now, let G^{-1} be the shadow of G . We can determine the similar weighting process on G^{-1} . Assume that the graph G is weighted by the above weighting process ω for the labeling set X . Define now the sets $-X$ and $-X_0$ by

$$-X \stackrel{def}{=} \{l_{-1}, \dots, l_{-N}\} \text{ and } -X_0 \stackrel{def}{=} \{l_0\} \cup (-X),$$

for all $j = 1, \dots, N$. Now, consider the set $V(G^{-1}) \times (-X) = V(G) \times (-X)$. Then, we can construct the weighting processes $(\omega_v^{-1})_{v \in V(G^{-1})}$ defined by

$$\omega_v^{-1} : E_{out}^v \rightarrow \{v\} \times (-X)$$

such that

$$\omega_v^{-1}(e^{-1}) = \omega_v^{-1}(e^{-1}v) \stackrel{def}{=} (v, l_{-i_j}),$$

whenever $w_v(e) = (v, l_{i_j}) \in \{v\} \times X$. Therefore, we can define the *weighting process ω^{-1} of $E(G^{-1})$* by

$$\omega^{-1} : E(G^{-1}) \rightarrow V(G^{-1}) \times (-X)$$

by

$$\omega^{-1}(e^{-1}) = \omega^{-1}(e^{-1}v) \stackrel{def}{=} (v, l_{-i_j}),$$

whenever

$$\omega(e) = (v, l_{ij}) \in \{v\} \times X.$$

Define the *canonical projection*

$$pr_G : V(G) \times X \rightarrow X$$

by

$$pr_G((v, l_k)) \stackrel{def}{=} l_k, \text{ for all } (v, l_k) \in V(G) \times X.$$

Similarly, define the canonical projection

$$pr_{G^{-1}} : V(G) \times (-X) \rightarrow -X$$

by

$$pr_{G^{-1}}((v, l_{-k})) \stackrel{def}{=} l_{-k}, \text{ for all } (v, l_{-k}) \in V(G^{-1}) \times (-X).$$

Definition 4.4. Let G be a locally finite connected directed graph equipped with its weighting process ω of the edge set $E(G)$ by $V(G) \times X$, where $X = \{l_1, \dots, l_N\}$, and $N = \max\{\deg_{out}(v) : v \in V(G)\}$. Sometimes, we denote the graph G equipped with the weighting process ω by the pair (G, ω) . Then the weighted graph (G, ω) is called the *canonical weighted graph*. Similarly, the weighted graph (G^{-1}, ω^{-1}) equipped with the weighting process ω^{-1} of $E(G^{-1})$ by $(-X) \times V(G)$ is called the *canonical weighted shadow* of (G, ω) , where G^{-1} is the shadow of G .

Let (G, ω) (resp. (G^{-1}, ω^{-1})) be the canonical weighted graph (resp. the canonical weighted shadow of (G, ω)). The image $pr_G \circ \omega$ of $E(G)$ (resp. $pr_{G^{-1}} \circ \omega^{-1}$), contained in X (resp. in $-X$), is called the *canonical labeling process* of (G, ω) (resp. of (G^{-1}, ω^{-1})).

Let (G, ω) be the canonical weighted graph with its canonical weighted shadow (G^{-1}, ω^{-1}) . Then we can extend this weighting processes ω and ω^{-1} to the weighting process, also denoted by ω , the free semigroupoid $\mathbb{F}^+(\widehat{G})$ of the shadowed graph $\widehat{G}$ of G . Let $\pm X_0^*$ be the collection of all finite words in $\pm X_0 = X_0 \cup (-X)$. It is easy to check that

$$\pm X_0^* = \mathcal{L}_N,$$

where $\mathcal{L}_N$ is the lattice path set generated by $\pm X = \{l_{\pm 1}, \dots, l_{\pm N}\}$, including l_0 .

Define a map

$$\omega : FP(\widehat{G}) \cup \{\emptyset\} \rightarrow V(G)^2 \times X^*$$

by

$$\omega(\emptyset) \stackrel{def}{=} (\emptyset, l_0)$$

and

$$\omega(w) = \omega(v \ e_1 \ \dots \ e_k v') \stackrel{\text{def}}{=} \left((v, \ v'), \ \prod_{j=1}^k \tilde{pr}(\omega_{\pm}(e_j)) \right),$$

where $\emptyset$ is the empty set in $V(G)^2$ and $FP(\widehat{G})$ is the (non-reduced) finite path set of the shadowed graph $\widehat{G}$, and where

$$\tilde{pr} \stackrel{\text{def}}{=} pr_G \cup pr_{G^{-1}} \text{ and } \omega_{\pm} \stackrel{\text{def}}{=} \omega \cup \omega^{-1},$$

whenever

$$w = v \ w \ v' = e_1 \ \dots \ e_k,$$

with

$$v, v' \in V(\widehat{G}), \text{ and } e_1, \dots, e_k \in E(\widehat{G}),$$

for $k \in \mathbb{N}$. i.e.,

$$\omega_{\pm}(e) = \begin{cases} \omega(e) & \text{if } e \in E(G) \\ \omega^{-1}(e) & \text{if } e \in E(G^{-1}), \end{cases}$$

and

$$\tilde{pr}(g) = \begin{cases} pr_G(g) & \text{if } g \in V(G) \times X \\ pr_{G^{-1}}(g) & \text{if } g \in V(G) \times (-X). \end{cases}$$

The above map ω from $FP(\widehat{G}) \cup \{\emptyset\}$ can be extended to $\mathbb{F}^+(\widehat{G})$ into $V(G)^2 \times (\pm X_0^*)$ by putting the weights on vertices:

$$\omega(v) \stackrel{\text{def}}{=} ((v, \ v), \ l_0), \text{ for all } v \in V(G).$$

Definition 4.5. *The above weighting process ω of the free semigroupoid $\mathbb{F}^+(\widehat{G})$ of $\widehat{G}$ is called the canonical weighting process of $\mathbb{F}^+(\widehat{G})$ or that of $\mathbb{G}$ induced by the canonical weighted graph (G, ω) .*

Remark 4.1. *Set-theoretically, the graph groupoid $\mathbb{G}$ is contained in the free semigroupoid $\mathbb{F}^+(\widehat{G})$ of the shadowed graph $\widehat{G}$ of G . So, we can determine the weighting process ω of $\mathbb{G}$ simply by restricting ω of $\mathbb{F}^+(\widehat{G})$. i.e., the weighting process ω of $\mathbb{G}$ is understood as $\omega = \omega|_{\mathbb{G}}$. We will use the notation ω for the weighting processes of $\mathbb{F}^+(\widehat{G})$ and $\mathbb{G}$, alternatively.*

Notice that the above weighting process ω of $\mathbb{F}^+(\widehat{G})$ represents the full admissibility conditions of $\mathbb{G}$. i.e., the weight $\omega(w)$ of $w \in \mathbb{G}$ contains its initial and terminal vertices and the connecting pattern (admissibility of edges) of w , in terms of labelings in $\pm X_0 = X_0 \cup (-X)$.

Definition 4.6. *Let (G, ω) be the canonical weighted graph with its canonical weighted shadow (G^{-1}, ω^{-1}) . Let $\mathbb{G}$ be the graph groupoid of G and assume that*

ω is the canonical weighting process of $\mathbb{G}$ induced by (G, ω) . Then we call the pair $(\mathbb{G}, \omega)$ the labeled graph groupoid of (G, ω) . We call the processes $\tilde{pr} \circ \omega$ on $(\mathbb{G}, \omega)$, the labeling process, where $\tilde{pr} = pr_G \cup pr_{G^{-1}}$.

Assumption From now, if we mention (locally finite connected) directed graphs, then they are automatically assumed to be the canonical weighted graphs. Similarly, if we mention graph groupoids, then they are also automatically assumed to be the labeled graph groupoids. $\square$


Example 4.1. Let G be the circulant graph with three vertices with

$$V(G) = \{v_1, v_2, v_3\}$$

and

$$E(G) = \{e_j = v_j \rightarrow v_{j+1} : j = 1, 2, 3, \text{ with } v_4 \stackrel{\text{def}}{=} v_1\}.$$

i.e.,



Then, since $\deg_{out}(v_j) = 1$, for all $j = 1, 2, 3$, we can get the labeling sets

$$X = \{l_1\} = \left\{ \overrightarrow{(1, e)} \right\} \text{ and } X_0 = \{l_0, l_1\},$$

contained in the lattice path set $\mathcal{L}_1$. Relatively, we have that

$$-X = \{l_{-1}\} = \left\{ \overrightarrow{(1, -e)} \right\},$$

for the shadow G^{-1} of G . Then we can get the weights

$$\omega(e_j) = ((v_j, v_{j+1}), l_1), \text{ for all } j = 1, 2, 3$$

and

$$\omega^{-1}(e_j^{-1}) = ((v_{j+1}, v_j), l_{-1}), \text{ for all } j = 1, 2, 3,$$

where $v_4 \stackrel{\text{def}}{=} v_1$, in $V(G)$. We can construct the labeled graph groupoid $(\mathbb{G}, \omega)$. Let $w = e_2 e_3 e_1 \in FPr(\widehat{G})$. Then we can get that

$$\omega(w) = ((v_2, v_2), l_1 l_1 l_1), \text{ and } \omega(w^{-1}) = ((v_2, v_2), l_{-1} l_{-1} l_{-1}).$$

Take now the reduced finite path $y = e_1^{-1} e_3^{-1} \in FPr(\widehat{G})$. Then we can get

$$\omega(y) = ((v_2, v_3), l_{-1}l_{-1}).$$

Suppose we have $w^2 = e_2 e_3 e_1 e_2 e_3 e_1$ in $FP_r(\widehat{G})$. Then the weight of w^2 is

$$\omega(w^2) = ((v_2, v_2), l_1 l_1 l_1 l_1 l_1 l_1).$$

4.2. The Operation θ and ω_+ .

In this section, we will define some operations on the labeling set $\pm X_0 = X_0 \cup (-X)$ of the labeled graph groupoid $(\mathbb{G}, \omega)$, induced by the canonical weighted graph (G, ω) . Then Define an operation

$$\theta : \pm X_0^* = \mathcal{L}_N \rightarrow \mathbb{Z}$$

by

$$\theta(l_{i_1} l_{i_2} \dots l_{i_n}) \stackrel{\text{def}}{=} \sum_{j=1}^n i_j,$$

for all $l_{i_1} \dots l_{i_n} \in \pm X_0^* = \mathcal{L}_N$, for $n \in \mathbb{N}$, where the addition of the right-hand side means the vector addition on $\mathbb{R}^2$.

By help of θ , we can define an operation,

$$\omega_+ : V(G)^2 \times (\pm X_0^*) \rightarrow V(G)^2 \times \mathbb{Z}$$

by

$$\begin{aligned} \omega_+((v, v'), l_{i_1} \dots l_{i_n}) &\stackrel{\text{def}}{=} ((v, v'), \theta(l_{i_1} \dots l_{i_n})) \\ &= \left((v, v'), \sum_{j=1}^n i_j \right), \end{aligned}$$

for all $((v, v'), l_{i_1} \dots l_{i_n}) \in V(G)^2 \times (\pm X_0^*)$. Later, we will use the above operation ω_+ to verify the amalgamated moments of the labeling operators on right graph von Neumann algebras. In conclusion, we can get that:

Proposition 4.2. *Let w_1 and w_2 be the reduced finite paths in $FP_r(\widehat{G})$ given in the previous paragraph. Then*

$$\theta((\tilde{p}r \circ \omega)(w_1 w_2)) = 0 \text{ in } \mathbb{Z},$$

and

$$\omega_+(\omega(w_1 w_2)) = ((v, v), 0),$$

if and only if the element $w_1 w_2 = v$ in $\mathbb{G} \setminus \{\emptyset\}$.

Proof. $(\Rightarrow)$ Suppose $\omega_+(\omega(w_1 w_2)) = ((v, v), 0)$. By hypothesis, we have that

$$\omega(w_1 w_2) = ((v, v), (l_{i_1} \dots l_{i_n} l_{j_1} \dots l_{j_k})),$$

whenever

$$\omega(w_1) = ((v, v'), l_{i_1} \dots l_{i_n})$$

and

$$\omega(w_2) = ((v', v), l_{j_1} \dots l_{j_k}).$$

By assumption, $\theta(l_{i_1} \dots l_{i_n} l_{j_1} \dots l_{j_k}) = 0$. And since w_1 and w_2 are “reduced” finite paths in $\mathbb{G}$, we can verify that (i) $n = k$, and (ii) for any l_{i_p} , there should be the unique entry $l_{j_q} = l_{-i_p}$, for $p, q = 1, \dots, n = k$. This shows that, if $w_1 = e_{i_1} \dots e_{i_n}$ and $w_2 = e_{j_1} \dots e_{j_n}$ in $FP_r(\widehat{G})$, with $e_{i_1}, \dots, e_{i_n}, e_{j_1}, \dots, e_{j_n} \in E(\widehat{G})$, then $e_{j_q} = e_{i_p}^{-1}$, uniquely, for $p, q = 1, \dots, n$. Therefore, $w_1 w_2 = v$.

$(\Leftarrow)$ Suppose now $w_1 w_2 = v \in V(G)$. Then

$$\omega_+(\omega(w_1 w_2)) = \omega_+(\omega(v)) = \omega_+((v, v), l_0) = ((v, v), 0).$$

■

The above proposition shows how the labeling on the graph groupoid $\mathbb{G}$ works. The labeling process on $\mathbb{G}$ gives a way to check how admissibility works under (RR).

4.3. Graph Automata.

In this section, we will construct an automaton induced by the given canonical weighted graph $G = (G, \omega)$. Let $X_0 = \{l_0, l_1, \dots, l_N\}$ be the labeling set of the countable locally finite connected directed graph G , consisting of lattices in $\mathbb{R}^2$, where

$$N \stackrel{def}{=} \max\{\deg_{out}(v) : v \in V(G)\},$$

and let $\pm X_0 = X_0 \cup (-X) = \{l_0, \pm l_1, \dots, \pm l_N\}$. Recall that the collection $\pm X_0^*$ of all words in $\pm X_0$ is identical to the lattice path set $\mathcal{L}_N$.

Under the above setting, we can create the automaton $\mathcal{A}_G = \langle \mathcal{X}_0, E(\widehat{G}), \varphi, \psi \rangle$, where $\widehat{G}$ is the shadowed graph of G and

$$\mathcal{X}_0 \stackrel{def}{=} \{\emptyset_G\} \cup (V(G)^2 \times (\pm X_0)),$$

and

$$\varphi(((v_1, v_2), l), e) \stackrel{\text{def}}{=} \begin{cases} \omega(e) & \text{if } e = v_2 e \text{ in } E(\widehat{G}), \text{ equivalently,} \\ & \exists v' \in V(G) \text{ and } l' \in \pm X_0 \text{ s.t.,} \\ & \omega(e) = ((v_2, v'), l') \neq \emptyset_G \\ \emptyset_G & \text{otherwise,} \end{cases}$$

and

$$\psi(((v_1, v_2), t), e) \stackrel{\text{def}}{=} \begin{cases} e & \text{if } e = v_2 e \text{ in } E(\widehat{G}) \\ \emptyset & \text{otherwise,} \end{cases}$$

for all $e \in E(\widehat{G})$, where $\emptyset_G = (\emptyset, l_0)$ is the *empty element* of $\mathcal{X}_0$ and $\emptyset$ is the empty element in $\mathbb{F}^+(\widehat{G})$.

Definition 4.7. We will say that the automaton $\mathcal{A}_G = \langle \mathcal{X}_0, E(\widehat{G}), \varphi, \psi \rangle$ is the graph automaton induced by the canonical weighted graph $G = (G, \omega)$. Sometimes, we call φ and ψ , the labeling map and the shift (or shifting map), respectively.

We can realize that the graph automaton $\mathcal{A}_G$ induced by G is identically same with the automaton $\mathcal{A}_{G^{-1}}$ induced by the shadow G^{-1} of G . Now, fix an edge $e_0 \in E(\widehat{G})$ and a reduced finite path $w = v_1 e_1 \dots e_n v'_n \in FP_r(\widehat{G})$, with $e_j = v_j e_j v'_j \in E(\widehat{G})$, where $v_j, v'_j \in V(\widehat{G})$, for all $j = 1, \dots, n$, for some $n \in \mathbb{N}$. Then we can extend φ and ψ inductively on the free semigroupoid $\mathbb{F}^+(\widehat{G})$ of $\widehat{G}$ as follows:

$$\begin{aligned} & \varphi(((v_1, v'_n), \omega(e_1) \dots \omega(e_n)), e_0) \\ &= \varphi(\omega(w), e_0) \\ & \stackrel{\text{def}}{=} \begin{cases} \omega(e_0) & \text{if } w e_0 \neq \emptyset \\ \emptyset_G & \text{otherwise,} \end{cases} \end{aligned}$$

and similarly,

$$\begin{aligned} & \psi(((v_1, v'_n), \omega(e_1) \dots \omega(e_n)), e_0) \\ &= \psi(\omega(w), e_0) \\ & \stackrel{\text{def}}{=} \begin{cases} e_0 & \text{if } w e_0 \neq \emptyset \\ \emptyset & \text{otherwise,} \end{cases} \end{aligned}$$

for $n \in \mathbb{N}$. Also, inductively, we have that, if $w' = e'_1 \dots e'_m \in FP_r(\widehat{G})$, for $m \in \mathbb{N}$, then

$$\begin{aligned} & \varphi(((v_1, v'_n), \omega(e_1) \dots \omega(e_n)), w') \\ & \stackrel{\text{def}}{=} \begin{cases} \omega(e'_m) & \text{if } w w' \neq \emptyset \\ \emptyset_G & \text{otherwise,} \end{cases} \end{aligned}$$

and similarly,

$$\psi(((v_1, v'_n), \omega(e_1) \dots \omega(e_n)), w')$$

$$\stackrel{def}{=} \begin{cases} w' & \text{if } w \text{ and } w' \neq \emptyset \\ \emptyset & \text{otherwise.} \end{cases}$$

Remark 4.2. As we have seen in the previous paragraphs, the (extended) definitions of φ and ψ from $\mathbb{F}^+(\widehat{G})$ into $V(G)^2 \times (\pm X_0^*)$ have the following meaning: by definition, we have $\varphi(\omega(w), w') = \omega(e')$, whenever w and $w' = e' \dots e''$ are admissible, where $\omega(w) \in V(G)^2 \times (\pm X_0^*)$ and $e', \dots, e'' \in E(\widehat{G})$, and hence it represents the admissibility of w and w' , in terms of the weighting process ω . Equivalently,

Computing $\varphi(\omega(w), w')$ means: if w and w' are admissible, what is the weight of the connecting edge e' of w' ?

Under the same setting, we have $\psi(\omega(w), w') = e'$: it again represents the admissibility of w and w' , in terms of the edges. Thus, roughly speaking, the labeling map φ and the shifting ψ represent the admissibility conditions on $\mathbb{F}^+(\widehat{G})$ (and hence $\mathbb{G}$). It gives more detailed information about $\mathbb{F}^+(\widehat{G})$ (or $\mathbb{G}$). Equivalently,

Computing $\psi(\omega(w), w')$ means: if w and w' are admissible, what is the starting edge of w' ?

Construct the automata actions $\{\mathcal{A}_e : e \in E(\widehat{G})\}$ of $\mathcal{A}_G$, acting on the set $V(G)^2 \times (\pm X_0^*)$ (consisting of all finite words in $\mathcal{X}_0$). Indeed, we can define the action $\mathcal{A}_e$ by

$$\mathcal{A}_e(((v_1, v_2), l_{i_1} \dots l_{i_n})) \stackrel{def}{=} \varphi(((v_1, v_2), l_{i_1} \dots l_{i_n}), e)$$

for all $((v_1, v_2), l_{i_1} \dots l_{i_n}) \in V(G)^2 \times (\pm X_0^*)$, for all $e \in E(\widehat{G})$. It is easy to be checked that the actions $\mathcal{A}_e$'s satisfies

$$\mathcal{A}_{e_1} \mathcal{A}_{e_2} = \mathcal{A}_{e_2 e_1}, \text{ for all } e_1, e_2 \in E(\widehat{G}).$$

Thus the actions $\{\mathcal{A}_v : v \in V(\widehat{G})\}$ are induced by $\mathcal{A}_e$'s since

$$\mathcal{A}_v \stackrel{def}{=} \mathcal{A}_{e e^{-1}} = \mathcal{A}_{e^{-1}} \mathcal{A}_e, \text{ whenever } e = v e \in E(\widehat{G}).$$

This guarantees that the automata actions $\{\mathcal{A}_e : e \in E(\widehat{G})\}$ on $V(G)^2 \times \pm X_0^*$ generate the groupoid which is groupoid-isomorphic to the graph groupoid $\mathbb{G}$ of G . i.e., we can have that:

Theorem 4.3. *The set $\{\mathcal{A}_e : e \in E(\widehat{G})\}$ of automata actions, acting on $V(G) \times (\pm X_0^*)$, generates the actions $\{\mathcal{A}_w : w \in \mathbb{F}^+(\widehat{G})\}$ of $\mathcal{A}_G$, acting on the same set. Moreover, the groupoid generated by this set $\{\mathcal{A}_e : e \in E(\widehat{G})\}$ is groupoid-isomorphic to the graph groupoid $\mathbb{G}$ of G . $\square$*

4.4. Graph-Automata Trees.

As before, the standing assumption for our graphs G are as follows: locally finite, connected, and countable. The edges of G are directed, and they are assigned weights with the use of a labeling set. In this section, we will consider graph-automata trees where the labeled graph groupoids act on. Let $\mathbb{G}$ be the labeled graph groupoid induced by the canonical weighted graph $G = (G, \omega)$, with the labeling set $\pm X_0 = \{l_0, \pm l_1, \dots, \pm l_N\}$, and let the canonical weighting process ω of the free semigroupoid $\mathbb{F}^+(\widehat{G})$ of the shadowed graph $\widehat{G}$ of G , be given as before, where

$$N \stackrel{def}{=} \max\{\deg_{out}(v) : v \in V(G)\}.$$

The canonical weighted graph G creates the corresponding graph automaton

$$\mathcal{A}_G = \langle \mathcal{X}_0, E(\widehat{G}), \varphi, \psi \rangle,$$

where

$$\mathcal{X}_0 = \{\emptyset_*\} \cup (V(G)^2 \times (\pm X_0)),$$

and φ is the labeling map and ψ is the shifting map. We already observed that $\mathbb{G}$ acts on a $E(\widehat{G})$ -set $V(G)^2 \times (\pm X_0^*)$. For convenience, we will denote the set $V(G)^2 \times (\pm X_0^*)$ by $\mathcal{X}_0^*$.

Recall that two countable directed graphs G_1 and G_2 are *graph-isomorphic*, if there exists a bijection

$$g : V(G_1) \cup E(G_1) \rightarrow V(G_2) \cup E(G_2),$$


such that (i) $g(V(G_1)) = V(G_2)$ and $g(E(G_1)) = E(G_2)$, and (ii) $g(e) = g(v_1 e v_2) = g(v_1) g(e) g(v_2)$ in $E(G_2)$, whenever $e = v_1 e v_2 \in E(G_1)$, with $v_1, v_2 \in V(G_1)$.

In [10] and [11], we showed that if two graphs G_1 and G_2 are graph-isomorphic, then the corresponding graph groupoids $\mathbb{G}_1$ and $\mathbb{G}_2$ are groupoid-isomorphic. More generally, if two graphs G_1 and G_2 have the graph-isomorphic shadowed graphs $\widehat{G}_1$ and $\widehat{G}_2$, then the graph groupoids $\mathbb{G}_1$ and $\mathbb{G}_2$ are groupoid-isomorphic. Also, we showed that if two graph groupoids $\mathbb{G}_1$ and $\mathbb{G}_2$ are groupoid-isomorphic, then the


(left) graph von Neumann algebras $\overline{\mathbb{C}[\mathbb{G}_1]}^w$ and $\overline{\mathbb{C}[\mathbb{G}_2]}^w$ are $*$ -isomorphic, as W^* -subalgebras in the operator algebra $B(H_{G_1}) = B(H_{G_2})$. Therefore, if two graphs G_1 and G_2 have the graph-isomorphic shadowed graphs, then the right graph von Neumann algebras M_{G_1} and M_{G_2} are $*$ -isomorphic, too.

Proposition 4.4. *Let G_1 and G_2 be countable directed graphs. If the shadowed graphs $\widehat{G}_1$ and $\widehat{G}_2$ are graph-isomorphic, then the right graph von Neumann algebras M_{G_1} and M_{G_2} are $*$ -isomorphic. $\square$*


Recall also that we say a directed graph G is a (directed) *tree*, if this graph G is connected and it has no loop finite paths in the free semigroupoid $\mathbb{F}^+(G)$ of G . Also, we say that a directed tree is *rooted*, if we can find-and-fix a vertex v_0 of G , with $\deg_{in}(v_0) = 0$. This fixed vertex v_0 is called the *root of G* . For instance, a graph



is a rooted tree with its (fixed) root v_0 . A rooted tree G is *one-flow*, if the directions of edges oriented only one way from the root v_0 . For example, a graph



is a one-flow rooted tree with its root v_0 . A one-flow rooted tree G is said to be *growing*, if G is an infinitely countable directed graph. Finally, we will say that a one-flow growing rooted tree G is *regular*, if the out-degrees of all vertices are identical. For example, a graph



is a regular one-flow growing rooted tree. In particular, if $\deg_{out}(v) = N$, for all vertices v , then this regular one-flow growing rooted tree is called *the N -regular tree*. The very above example is the 2-regular tree.

Let N be the maximal out-degree of the graph G , and let $\mathcal{T}_{2N}$ be the $2N$ -regular tree. Then the automata actions $\{\mathcal{A}_w : w \in \mathbb{F}^+(\widehat{G})\}$ of the graph automaton $\mathcal{A}_G$ acts on this $2N$ -regular tree $\mathcal{T}_{2N}$. Indeed, for the set $\{\mathcal{A}_w\}_{w \in \mathbb{F}^+(\widehat{G})}$ of automata actions of the graph automaton $\mathcal{A}_G$, we can create a one-flow growing rooted tree $\mathcal{T}_G$ having its “arbitrarily” fixed root $\omega(v_0) = ((v_0, v_0), x_0) \in \mathcal{X}_0^*$, where

$$V(\mathcal{T}_G) \stackrel{def}{=} \left\{ \varphi(X, e) \mid X \in \mathcal{X}_0^*, e \in E(\widehat{G}) \right\}$$

and

$$E(\mathcal{T}_G) \stackrel{def}{=} \left\{ \psi(X, e) \mid X \in \mathcal{X}_0^*, e \in E(\widehat{G}) \right\},$$

where $\widehat{G}$ is the shadowed graph of G .

Remark 4.3. Notice that, by the connectedness of the shadowed graph $\widehat{G}$ of the connected graph G , we can fix any weight $\omega(v)$ of $v \in V(\widehat{G})$, as the root of the tree $\mathcal{T}_G$. Suppose $\mathcal{T}_{v_1}$ and $\mathcal{T}_{v_2}$ are the one-flow growing trees with their roots $((v_1, v_1), x_0)$ and $((v_2, v_2), x_0)$, respectively. Then, $\mathcal{T}_{v_i}$ is embedded in $\mathcal{T}_{v_j}$, as a full-subgraph (See below), whenever $i \neq j \in \{1, 2\}$. In general, the graphs $\mathcal{T}_{v_1}$ and $\mathcal{T}_{v_2}$ have no graph-isomorphic relation, but they are embedded from each other, by the connectedness of $\widehat{G}$. Since v_1 and v_2 are arbitrary, we can consider only one choice $\mathcal{T}_v$, for $v \in V(G)$, as a candidate of the tree, where $\{\mathcal{A}_w\}_{w \in \mathbb{F}^+(\widehat{G})}$ act. Denote it by $\mathcal{T}_G$. i.e., whenever we choose one tree $\mathcal{T}_v$, for $v \in V(G)$, then the trees $\mathcal{T}_{v'}$ ’s are embedded in $\mathcal{T}_v$, for all $v' \in V(G)$. Without loss of generality, if we write $\mathcal{T}_G$ from now, then it means a tree $\mathcal{T}_v$, for a fixed vertex $v \in V(G)$. Remark that, the tree $\mathcal{T}_G$ has its root $\omega(v)$, if and only if $\mathcal{T}_G = \mathcal{T}_v$.

We can easily check that

$$FP(\mathcal{T}_G) = \left\{ \psi(X, w) \mid X \in \mathcal{X}_0^*, w \in FP(\widehat{G}) \right\},$$

and hence

$$\begin{aligned} \mathbb{F}^+(\mathcal{T}_G) = & \{ \emptyset_G \} \cup \left\{ \varphi(X, e) \mid \begin{array}{l} X \in \mathcal{X}_0^*, \\ e \in E(\widehat{G}) \end{array} \right\} \\ & \cup \left\{ \psi(X, w) \mid \begin{array}{l} X \in \mathcal{X}_0^*, \\ w \in FP(\widehat{G}) \end{array} \right\}. \end{aligned}$$

By the connectedness of the graph G , and by the definition of the automata actions, every (nonempty) finite paths in $FP(\widehat{G})$ is embedded in the tree $\mathcal{T}_G$, via the automata actions. Then, we can construct the full-subgraphs $\{\mathcal{T}_w : w \in FP(\widehat{G})\}$ of $\mathcal{T}_G$, where $\mathcal{T}_w$'s are the one-flow growing rooted tree with their roots $\omega(w)$, for all $w \in FP(\widehat{G})$.

Recall that, we say that a countable directed graph G_1 is a *full-subgraph* of a countable directed graph G_2 , if

$$E(G_1) \subseteq E(G_2)$$

and

$$V(G_1) = \left\{ v \in V(G_2) \mid \begin{array}{l} e = ve \text{ or } e = ev, \\ \forall e \in E(G_1) \end{array} \right\}.$$

Notice the difference between full-subgraphs and subgraphs: we say that G_1 is a *subgraph* of G_2 , if

$$V(G_1) \subseteq V(G_2)$$

and

$$E(G_1) = \left\{ e \in E(G_2) \mid \begin{array}{l} e = v e v', \\ \forall v, v' \in V(G_2) \end{array} \right\}.$$

Every subgraph is a full-subgraph, but the converse does not hold true, in general.

Definition 4.8. Let $\mathcal{T}_G$ be the above one-flow growing rooted tree, where the automata actions $\{\mathcal{A}_w : w \in \mathbb{F}^+(\widehat{G})\}$ of the graph automaton $\mathcal{A}_G$ act. This tree $\mathcal{T}_G$ is called the $\mathcal{A}_G$ -tree, which is a full-subgraph of the $2N$ -regular tree $\mathcal{T}_{2N}$. The full-subgraphs $\mathcal{T}_w$'s for $w \in FP(\widehat{G})$ of $\mathcal{T}_G$ are called the w -parts of $\mathcal{T}_G$.

The important thing is now that the w -parts $\mathcal{T}_w$'s of the $\mathcal{A}_G$ -graph $\mathcal{T}_G$ are embedded in the $2N$ -regular tree $\mathcal{T}_{2N}$, and $\mathcal{T}_{w'}$'s are embedded in $\mathcal{T}_w$'s, whenever

$$\varphi(\omega(v), w') = ((v, v'), X'),$$

where

$$\varphi(\omega(v), w) = ((v, v''), X) \text{ and } X' = (X, X''),$$

for some $X'' \in (\pm X_0^*)$, where $\omega(v)$ is the root of $\mathcal{T}_G = \mathcal{T}_v$.

Remark 4.4. *The construction of the $\mathcal{A}_G$ -tree $\mathcal{T}_G$ is nothing but the rearrangement of the finite paths in the free semigroupoid $\mathbb{F}^+(\widehat{G})$ of the shadowed graph $\widehat{G}$ inside the $2N$ -regular tree $\mathcal{T}_{2N}$, up to the admissibility on $\mathbb{F}^+(\widehat{G})$. Notice that, in fact, the $\mathcal{A}_G$ -tree $\mathcal{T}_G$ contains the information about the vertices in $\mathbb{F}^+(\widehat{G})$, too, since the vertices of $\mathcal{T}_G$ are contained in $\mathcal{X}_0^* = V(G)^2 \times (\pm X_0^*)$. Remark that, by the connectedness of $\widehat{G}$, the w -parts $\mathcal{T}_w$'s of $\mathcal{T}_G$ for $w \in FP(\widehat{G})$ are well-constructed as a one-flow growing tree with their roots $\varphi(\omega(v), w)$, where $\omega(v)$ is the root of $\mathcal{T}_G$. Moreover, each tree $\mathcal{T}_w$ is embedded in the other trees $\mathcal{T}_G$.*

Remark that, even though $|\mathbb{G}| < \infty$, in general, $|\mathbb{F}^+(\widehat{G})| = \infty$, whenever $|E(G)| \geq 1$. In [17], we observed the following special labeled graph groupoids.

Definition 4.9. *Let G , $\mathbb{F}^+(\widehat{G})$, and $\mathbb{G}$ be given as before and let $\mathcal{A}_G = \langle \mathcal{X}_0, E(\widehat{G}), \varphi, \psi \rangle$ be the graph automaton induced by G , acting on the $2N$ -regular tree $\mathcal{T}_{2N}$. For any fixed $w \in FP(\widehat{G})$, the tree $\mathcal{T}_w$ is the w -part of the $\mathcal{A}_G$ -tree $\mathcal{T}_G$, with its root $\varphi(\omega(v), w)$, where $\omega(v)$ is the fixed root of the $\mathcal{A}_G$ -tree $\mathcal{T}_G$. Let $\mathbb{G}(\mathcal{T}_w)$ be the groupoid generated by the actions $\mathcal{A}_y$'s acting only on $\mathcal{T}_w$. If $\mathbb{G}(\mathcal{T}_w)$'s are groupoid-isomorphic to $\mathbb{G}(\mathcal{T}_G)$, for all $w \in FP(\widehat{G})$, then we say that the groupoid $\mathbb{G}(\mathcal{A}_G)$, generated by $\mathcal{A}_G$, is a fractaloid. Equivalently, we say that the (labeled) graph groupoid $\mathbb{G}$ is a fractaloid.*

Notice that

$$\mathbb{G}(\mathcal{A}_G) \stackrel{\text{Groupoid}}{=} \mathbb{G} \stackrel{\text{Groupoid}}{=} \mathbb{G}(\mathcal{T}_G).$$

The first groupoid-isomorphic relation is shown in the previous section, and the second groupoid-isomorphic relation holds, by the previous remark. Readers can understand the above definition of fractaloids as the graph-groupoid version of the fractal groups (See [1] and [17]). The following theorem provides the graph-theoretical characterization of fractaloids.

Theorem 4.5. *(See [17]) Let G be a canonical weighted graph with its labeled graph groupoid $\mathbb{G}$, and let $\mathcal{A}_G$ be the graph automaton induced by G and $\mathcal{T}_G$, the $\mathcal{A}_G$ -tree. Every w -part $\mathcal{T}_w$ of $\mathcal{T}_G$ is graph-isomorphic to $\mathcal{T}_G$, for all $w \in FP(\widehat{G})$, if and only if $\mathbb{G}$ is a fractaloid. $\square$*

Let G be a canonical weighted graph with its labeled graph groupoid $\mathbb{G}$, and assume that the automata actions $\{\mathcal{A}_w : w \in FP(\widehat{G})\}$ of the graph automaton $\mathcal{A}_G$ act fully on the $2N$ -regular tree $\mathcal{T}_{2N}$, in the sense that the $\mathcal{A}_G$ -tree $\mathcal{T}_G$, which is a full-subgraph of $\mathcal{T}_{2N}$, is graph-isomorphic to $\mathcal{T}_{2N}$. i.e., the automata actions of $\mathcal{A}_G$ act fully on $\mathcal{T}_{2N}$, if $\mathcal{T}_G \stackrel{\text{Graph}}{=} \mathcal{T}_{2N}$.

Corollary 4.6. *(See [17]) Let G be a canonical weighted graph with its labeled graph groupoid $\mathbb{G}$, and let $\mathcal{A}_G$ be the graph automaton induced by G . The automata*

actions $\{\mathcal{A}_w : w \in FP(\widehat{G})\}$ of $\mathcal{A}_G$ act fully on the $2N$ -regular tree $\mathcal{T}_{2N}$, if and only if $\mathbb{G}$ is a fractaloid. $\square$

The above theorem provides the graph-theoretical characterization of fractaloids: $\mathbb{G}$ is a fractaloid if and only if

$$\mathcal{T}_G \stackrel{\text{Graph}}{=} \mathcal{T}_{2N} \stackrel{\text{Graph}}{=} \mathcal{T}_w, \text{ for all } w \in \mathbb{F}^+(\widehat{G}).$$

And the above corollary provides the algebraic (or automata-theoretical) characterization of fractaloids; $\mathbb{G}$ is a fractaloid if and only if the automata actions of the graph automaton $\mathcal{A}_G$ act fully on $\mathcal{T}_{2N}$.

5. LABELING OPERATORS OF GRAPH GROUPOIDS

In this section, we define labeling operators of labeled graph groupoids in right graph von Neumann algebras. Let G be a canonical weighted graph with its labeled graph groupoid $\mathbb{G}$. Then the labeling process on $\mathbb{G}$ induces the labeling operator T_G of $\mathbb{G}$ contained in the right graph von Neumann algebra $M_G = \overline{\mathbb{C}[\mathbb{G}]}^w$ of G , in $B(H_G)$.

5.1. The Labeling Operator of $\mathbb{G}$ in M_G .

Let $G = (G, \omega)$ be a canonical weighted graph, where G is a countable locally finite connected directed graph, and let $\mathbb{G} = (\mathbb{G}, \omega)$ be the corresponding labeling graph groupoid of G . Let H_G be the graph Hilbert space in the sense of Section 2.3. Throughout this section, put

$$N = \max\{\deg_{out}(v) : v \in V(G)\}.$$

Definition 5.1. Define an operator $T_k \in B(H_G)$ by

$$T_k \xi_w \stackrel{\text{def}}{=} \xi_w \xi_{e_{l_k}} = \xi_{we_{l_k}}$$

$$= \begin{cases} \xi_{we_k} & \text{if } \exists e_{l_k} \in E(\widehat{G}) \text{ s.t.} \\ & w e_{l_k} \neq \emptyset \text{ in } FP_r(\widehat{G}) \\ & \text{and } (\tilde{p}r \circ \omega)(e_{l_k}) = l_k \\ \xi_\emptyset = 0_{H_G} & \text{otherwise,} \end{cases}$$

for all $w \in \mathbb{G}$, where $\xi_w \in \mathcal{B}_{H_G} \cup \{\xi_v : v \in V(\widehat{G})\}$, and where $(\tilde{p}\tilde{r} \circ \omega)(e_{l_k}) = l_k \in \pm X_0$ is the weight of the edge e_{l_k} , where $\pm X_0 = \{l_0, \pm l_1, \dots, \pm l_N\}$, for $k = \pm 1, \dots, \pm N$. These operators T_k 's are called the k -th labeling operators, for all $k = \pm 1, \dots, \pm N$. Then we can define the operator $T_G \in B(H_G)$ by

$$T_G \stackrel{\text{def}}{=} \sum_{k=-N}^{-1} T_k + \sum_{j=1}^N T_j.$$

This operator T is called the labeling operator of $\mathbb{G}$ on H_G . For convenience, denote $\sum_{k=-N}^{-1} T_k$ and $\sum_{j=1}^N T_j$ by T_- and T_+ , respectively.

The labeling operator T_G of $\mathbb{G}$ on H_G is a Heck-type operator or a Ruelle-operator-like operator (See [1], [17], [33] and [34]). The labeling operators are originally defined in [17]. The following proposition was already proven in [17]. It shows that we can regard the k -th labeling operators T_k 's, for $k = \pm 1, \dots, \pm N$, and the labeling operator T_G , as elements of the right graph von Neumann algebra M_G of G .

Proposition 5.1. (See [17]) *Let G be the given graph with its labeled graph groupoid $\mathbb{G}$, and let $M_G = \overline{\mathbb{C}[\mathbb{G}]}^w$ be the right graph von Neumann algebra of G , in $B(H_G)$. Then the k -th labeling operators T_k and the labeling operator T_G of $\mathbb{G}$ are contained in M_G , for $k = \pm 1, \dots, \pm N$. $\square$*

Indeed, we can define the elements $\tau_k \in M_G$, by

$$\tau_k \stackrel{\text{def}}{=} \sum_{e \in E(\widehat{G}), (\tilde{p}\tilde{r} \circ \omega)(e) = l_k} R_e \in M_G, \text{ for } k = \pm 1, \dots, \pm N.$$

Then these operator τ_k satisfies that

$$\begin{aligned} \tau_k \xi_w &= \sum_{e \in E(\widehat{G}), (\tilde{p}\tilde{r} \circ \omega)(e) = l_k} (R_e \xi_w) = \sum_{e \in E(\widehat{G}), (\tilde{p}\tilde{r} \circ \omega)(e) = l_k} \xi_{we} \\ &= \begin{cases} \xi_{we_0} & \text{if } we_0 \neq \emptyset \\ 0_{H_G} & \text{otherwise.} \end{cases} \end{aligned}$$

By the definition of labeling processes, (i) if the above H_G -value is nonzero, then there exists a “unique” $e_0 \in E(\widehat{G})$, with $\omega(e_0) = ((v, v'), l_k)$, whenever $w = wv$ and $e_0 = v e_0 v'$, and hence $(\tilde{p}\tilde{r} \circ \omega) = l_k$, and (ii) if the above H_G -value is zero, then we can conclude $w = wv$, with $\deg_{\text{out}}(v) = n_v$, and $|n_v| \not\geq |k|$, for $k \in \{\pm 1, \dots, \pm N\}$. i.e.,

$$\begin{cases} \xi_{we_0} = T_k \xi_w & \text{if } \exists! e_0 \in E(\widehat{G}) \text{ s.t.} \\ & (\tilde{p}r \circ \omega)(e_0) = x_k \\ \xi_\emptyset = 0_{H_G} & \text{otherwise.} \end{cases}$$

Therefore, the k -th labeling operators T_k are identical to $\tau_k \in M_G$, and hence, by definition, the labeling operator T_G of $\mathbb{G}$ is identified with the element $\tau \in M_G$, defined by

$$\tau = \left(\sum_{k=-N}^{-1} \tau_k \right) + \left(\sum_{j=1}^N \tau_j \right).$$

The above proposition shows that, without loss of generality, we can regard the k -th labeling operators T_k 's and the labeling operator T_G of $\mathbb{G}$ as elements of the right graph von Neumann algebra M_G , for $k = \pm 1, \dots, \pm N$. Recall that, the right graph von Neumann algebra $\mathbb{M}_G = \overline{\mathbb{C}[\mathbb{G}]}^w \subseteq B(H_G)$ is $*$ -isomorphic to a D_G -valued reduced free product algebra, where D_G is the $\mathbb{C}$ -diagonal subalgebra of M_G . Therefore, we can understand the operators T_k 's and T_G , as D_G -valued random variables in (M_G, E) , where $E : M_G \rightarrow D_G$ is the canonical conditional expectation. So, the labeling operators have their D_G -valued free distributional data.

5.2. D_G -Valued Free Distributional Data of Labeling Operators.

As before, let $G = (G, \omega)$ be a canonical weighted graph with its labeled graph groupoid $\mathbb{G} = (\mathbb{G}, \omega)$. In this section, we will consider the D_G -valued free distributional data of the k -th labeling operators T_k 's and the labeling operator T_G of $\mathbb{G}$ on the graph Hilbert space H_G , by regarding them as D_G -valued random variables in the right graph W^* -probability space (M_G, E) over the $\mathbb{C}$ -diagonal subalgebra D_G , where

$$N \stackrel{\text{def}}{=} \max\{\deg_{out}(v) : v \in V(G)\} \in \mathbb{N}.$$

Notation Without loss of generality, we will use the notations T_k 's and τ_k 's (resp., T_G and τ), alternatively, for $k = \pm 1, \dots, \pm N$. $\square$

Now, fix $k \in \{\pm 1, \dots, \pm N\}$, and consider the D_G -valued joint moments $E(T_k^{r_1} \dots T_k^{r_n})$ of the k -th labeling operator

$$T_k = \sum_{e \in E(\widehat{G}), (\tilde{p}r \circ \omega)(e) = l_k} R_e \in M_G,$$

where $(r_1, \dots, r_n) \in \{1, *\}^n$, for all $n \in \mathbb{N}$.

Lemma 5.2. (See [17]) Let T_k be the labeling operator on the graph Hilbert space H_G . Then its adjoint T_k^* is identified with the $(-k)$ -th labeling operator T_{-k} , for $k \in \{\pm 1, \dots, \pm N\}$. $\square$

Indeed, for each k , we can have

(5.1)

$$\begin{aligned} T_k^* &= \sum_{e \in E(\widehat{G}), (\widetilde{pr} \circ \omega)(e) = l_k} R_e^* = \sum_{e \in E(\widehat{G}), (\widetilde{pr} \circ \omega)(e) = l_k} R_{e^{-1}} \\ &= \sum_{e^{-1} \in E(\widehat{G}), (\widetilde{pr} \circ \omega)(e^{-1}) = l_{-k}} R_{e^{-1}} = T_{-k}, \end{aligned}$$

by the canonical weighting process, for $k = \pm 1, \dots, \pm N$. Thus, for any k , the adjoint T_k^* of the k -th labeling operator T_k is the $(-k)$ -th labeling operator T_{-k} , for all $k = \pm 1, \dots, \pm N$. By the above lemma, we can conclude the self-adjointness of the labeling operator T_G on H_G .

Corollary 5.3. (See [17]) Let T_G be the labeling operator of $\mathbb{G}$ on H_G . Then it is self-adjoint. $\square$

The above corollary guarantees that the study of D_G -valued free distributional data of T_G is to study the spectral property of T_G . Now, we will consider the D_G -freeness of $\{T_k, T_{-k}\}$'s in the graph W^* -probability space (M_G, E) , for all $k = 1, \dots, N$.

Theorem 5.4. (See [17]) Let $\{T_k : k = \pm 1, \dots, \pm N\}$ be the k -th labeling operators. Then the families $\{T_k, T_{-k}\}$'s are free over D_G from each other in (M_G, E) , for $k = 1, \dots, N$. i.e., the families $\{T_{k_1}, T_{-k_1}\}$ and $\{T_{k_2}, T_{-k_2}\}$ are free over $\mathbb{D}_G$ in $(\mathbb{M}_G, E)$, whenever $k_1 \neq k_2$ in $\{1, \dots, N\}$. $\square$

The above theorem can be proven, by the fact: the subsets

$$\mathcal{R}_k \stackrel{\text{def}}{=} \{e \in E(\widehat{G}) : (\widetilde{pr} \circ \omega)(e) = k \text{ or } -k\} \subset \mathbb{G},$$

for $k = 1, \dots, N$, are diagram-distinct from each other.

Consider now the D_G -valued joint $*$ -moments of $T_{\pm 1}, \dots, T_{\pm N}$. Let $(i_1, \dots, i_n) \in \{\pm 1, \dots, \pm N\}^n$, for $n \in \mathbb{N}$. Observe that

$$\begin{aligned} &E(T_{i_1} \dots T_{i_n}) \\ &= E \left(\left(\sum_{e_1 \in E(\widehat{G}), (\widetilde{pr} \circ \omega)(e_1) = l_{i_1}} R_{e_1} \right) \dots \left(\sum_{e_n \in E(\widehat{G}), (\widetilde{pr} \circ \omega)(e_n) = l_{i_n}} R_{e_n} \right) \right) \end{aligned}$$

$$\begin{aligned}
 &= E \left(\sum_{(e_1, \dots, e_n) \in E(\widehat{G})^n, (\widetilde{pr} \circ \omega)(e_j) = l_{i_j}} R_{e_n e_{n-1} \dots e_2 e_1} \right) \\
 (5.2) \quad &= E \left(\sum_{w \in E(\widehat{G})^n \cap FP(\widehat{G}), (\widetilde{pr} \circ \omega)(w) = l_{i_1} \dots l_{i_n}} R_w \right),
 \end{aligned}$$

where $E(\widehat{G})^n = \underbrace{E(\widehat{G}) \times \dots \times E(\widehat{G})}_{n\text{-times}}$, for $n \in \mathbb{N}$.

Now, recall the definition of the operations θ and ω_+ , introduced in Section 4.2.

Lemma 5.5. *Let $w \in E(\widehat{G})^n \cap FP(\widehat{G})$, with $\omega(w) = ((v, v), l_{i_1} \dots l_{i_n})$, for $n \in \mathbb{N}$, for some $v \in V(\widehat{G})$. Then $\omega_+(\omega(w)) = ((v, v), 0)$ if and only if $R_w = R_v$, for $v \in V(G)$.*

Proof. ($\Rightarrow$) Assume that $w \in \mathbb{G}$, and let

$$\begin{aligned}
 \omega_+(\omega(w)) &= \omega_+(((v, v), l_{i_1} \dots l_{i_n})) = ((v, v), \theta(l_{i_1} \dots l_{i_n})) \\
 &= ((v, v), \sum_{j=1}^n i_j) = ((v, v), 0).
 \end{aligned}$$

Then, we can conclude that $w = v w v$, for $v \in V(G)$, and $w = e_{i_1} \dots e_{i_n}$, with $e_{i_j} \in E(\widehat{G})$, having $\omega(e_{i_j}) = l_{i_j}$, for all $j = 1, \dots, n$. Since $\sum_{j=1}^n i_j = 0$ in $\mathbb{Z}$, the number n should be an even number in $2\mathbb{N}$, and $\{i_1, \dots, i_n\}$ is decomposed by $\{j_1, \dots, j_{\frac{n}{2}}\}$ and $\{q_1, \dots, q_{\frac{n}{2}}\}$. Furthermore, the set $\{e_{i_1}, \dots, e_{i_n}\}$ is decomposed by $\mathcal{E}_1 = \{e_{j_1}, \dots, e_{j_{\frac{n}{2}}}\}$ and $\mathcal{E}_2 = \{e_{q_1}, \dots, e_{q_{\frac{n}{2}}}\}$. By the definitions of ω_+ , we can conclude that $\mathcal{E}_2 = \mathcal{E}_1^{-1}$ in $E(\widehat{G})$.

Therefore, by [10] and [11], $w = v w v = v v v = v$ in $V(G)$.

($\Leftarrow$) Suppose $R_w = R_v$, for $v \in V(G)$. Then

$$\omega_+(\omega(v)) = \omega_+(((v, v), l_0)) = ((v, v), 0).$$

■

By the previous observations, we obtain the following theorem providing the D_G -valued free distributional data of the k -th labeling operators $T_{\pm 1}, \dots, T_{\pm N}$.

Theorem 5.6. *Let T_k 's be the k -th labeling operators on H_G , for all $k = \pm 1, \dots, \pm N$. Then the $\mathbb{D}_G$ -valued joint $*$ -moments of them is determined by*

(5.3)

$$E(T_{i_1} \dots T_{i_n}) = \sum_{w \in E(\widehat{G})^n \cap FP(\widehat{G}), \omega_+(\omega(w)) = ((v, v), 0)} R_w,$$

in $\mathbb{D}_G$, for all $(i_1, \dots, i_n) \in \{\pm 1, \dots, \pm N\}$, for $n \in \mathbb{N}$. $\square$

The formula (5.3) provides the D_G -valued free distributional data of $\{T_k : k = \pm 1, \dots, \pm N\}$ in (M_G, E) , in terms of their D_G -valued joint $*$ -moments. By the Moebius inversion (e.g., [5] and [21]: also see Section 1.1), we can get that the D_G -valued joint $*$ -cumulants of $\{T_k : k = \pm 1, \dots, \pm N\}$:

$$k_n(T_{i_1}, \dots, T_{i_n}) = \sum_{\pi \in NC(n)} E_\pi(T_{i_1}, \dots, T_{i_n}) \mu(\pi, 1_n).$$

Similar to [10] and [11], we obtain the following lemma.

Lemma 5.7. *Let $w_1, \dots, w_n \in \mathbb{G}$ and let $R_{w_1}, \dots, R_{w_n} \in M_G$ be the corresponding right multiplication operators, for $n \in \mathbb{N}$. Then*

(5.4)

$$k_n(R_{w_1}, R_{w_2}, \dots, R_{w_n}) = \mu_0 E(R_{w_n \dots w_2 w_1}),$$

where

$$\mu_0 = \sum_{\pi \in NC(w_1, \dots, w_n)} \mu(\pi, 1_n),$$

where

$$NC(w_1, \dots, w_n) \stackrel{\text{def}}{=} \left\{ \pi \in NC(n) \mid \begin{array}{l} E_\pi(R_{w_1}, \dots, R_{w_n}) \\ = E(R_{w_n \dots w_2 w_1}) \neq 0_{D_G} \end{array} \right\}.$$

$\square$

So, the above lemma will let us find the D_G -valued free distributional data of the k -th labeling operators $\{T_{\pm 1}, \dots, T_{\pm N}\}$, in terms of the D_G -valued joint $*$ -cumulants of them. Notice that, again by Moebius inversion, the D_G -valued free distributional data of $T_{\pm 1}, \dots, T_{\pm N}$, represented by D_G -valued $*$ -moments and by D_G -valued $*$ -cumulants, are equivalent.

Let $(i_1, \dots, i_n) \in \{\pm 1, \dots, \pm N\}^n$, for $n \in \mathbb{N}$. Observe that

$$\begin{aligned} & k_n(T_{i_1}, \dots, T_{i_n}) \\ &= k_n \left(\left(\sum_{e_1 \in E(\widehat{G}), (\widetilde{p\overline{r}} \circ \omega)(e) = l_{i_1}} R_{e_1} \right), \dots, \left(\sum_{e_n \in E(\widehat{G}), (\widetilde{p\overline{r}} \circ \omega)(e) = l_{i_n}} R_{e_n} \right) \right) \\ &= \sum_{(e_1, \dots, e_n) \in E(\widehat{G}), (\widetilde{p\overline{r}} \circ \omega)(e_{i_j}) = l_{i_j}} k_n(R_{e_1}, \dots, R_{e_n}) \end{aligned}$$

by the bimodule map property of $k_n(\dots)$ (See [21])

$$(5.5) \quad = \sum_{(e_1, \dots, e_n) \in E(\widehat{G})^n \cap FFP(\widehat{G}), (\widetilde{pr} \circ \omega)(e_{i_j}) = l_{i_j}} \left(\mu_{(e_1, \dots, e_n)} E(R_{e_n \dots e_2 e_1}) \right)$$

by (5.4), where

$$\mu_{(e_1, \dots, e_n)} = \sum_{\pi \in NC(e_1, \dots, e_n)} \mu(\pi, 1_n),$$

where

$$NC(e_1, \dots, e_n) = \left\{ \pi \in NC(n) \left| \begin{array}{l} E_\pi(R_{e_1}, \dots, R_{e_n}) \\ = E(R_{e_n e_{n-1} \dots e_2 e_1}) \\ \neq 0_{D_G} \end{array} \right. \right\}.$$

Thus, the formula (5.5) is identified with the following formula (5.6)

$$(5.6) \quad \sum_{w=e_1 \dots e_n \in E(\widehat{G})^n \cap FFP(\widehat{G}), \omega_+(\omega(w))=((v, v), 0)} \mu_w R_w,$$

where

$$(5.7) \quad \mu_w = \mu_{(e_1, \dots, e_n)}, \text{ whenever } w = e_n \dots e_1 \in \mathbb{G}.$$

Therefore, we can get the following equivalent D_G -valued free distributional data of the k -th labeling operators T_k 's, for $k = \pm 1, \dots, \pm N$:

Theorem 5.8. *Let T_k 's be the k -th labeling operators, for $k = \pm 1, \dots, \pm N$. Then the $\mathbb{D}_G$ -valued joint $*$ -cumulants of them are*

$$(5.8) \quad k_n(T_{i_1}, \dots, T_{i_n}) = \sum_{w \in E(\widehat{G})^n \cap FFP(\widehat{G}), \omega_+(\omega(w))=((v, v), 0)} (\mu_w R_w),$$

for all $(i_1, \dots, i_n) \in \{\pm 1, \dots, \pm N\}^n$, for $n \in \mathbb{N}$, where μ_w 's are given in (5.7). $\square$

In [10] and [11], we showed that $\mu_w = 0$, whenever there exists at least one pair $(i, j) \in \{1, \dots, n\}$ such that e_i and e_j are diagram-distinct, where $w = e_1 \dots e_n \in FFP(\widehat{G})$, with $e_1, \dots, e_n \in E(\widehat{G})$. Therefore, we can have the following corollary.

Corollary 5.9. *Let T_k 's be the k -th labeling operators, for $k = \pm 1, \dots, \pm N$. Then*

(1) $k_n(T_{i_1}, \dots, T_{i_n}) = 0_{D_G}$, whenever n is odd.

(2) $k_n(T_{i_1}, \dots, T_{i_n}) = \sum_{w \in \mathcal{W}_n^{(c)}(i_1, \dots, i_n)} (\mu_w R_w)$, where

(5.9)

$$\mathcal{W}_n^{(c)}(i_1, \dots, i_n) = \left\{ w = e_1 \dots e_n \left| \begin{array}{l} w \in E(\widehat{G})^n \cap FP(\widehat{G}) \\ (\tilde{p}r \circ \omega)(w) = l_{i_1} \dots l_{i_n} \\ \theta(l_{i_1} \dots l_{i_n}) = 0, \text{ and} \\ \exists e \in E(\widehat{G}) \text{ s.t. } e_1 \dots e_n \text{ is a} \\ \text{word only in } \{e, e^{-1}\}. \end{array} \right. \right\},$$

for all $n \in \mathbb{N}$. $\square$

Now, let's consider the D_G -valued free distributional data of the labeling operator T_G of $\mathbb{G}$ on the graph Hilbert space H_G . Recall that the labeling operator T_G is self-adjoint in $M_G \subseteq B(H_G)$. Therefore, it is enough to consider its D_G -valued (non-joint and non-*)moments $\{E(T_G^n)\}_{n \in \mathbb{N}}$. Since T_G is self-adjoint, the data $\{E(T_G^n) : n \in \mathbb{N}\}$ contain the full D_G -valued free distributional data of it.

Fix $n \in \mathbb{N}$ and consider T_G^n :

$$\begin{aligned} T_G^n &= (T_- + T_+)^n = (T_{-N} + \dots + T_{-1} + T_1 + \dots + T_N)^n \\ &= \sum_{(i_1, \dots, i_n) \in \{\pm 1, \dots, \pm N\}^n} T_{i_1} \dots T_{i_n}. \end{aligned}$$

Thus, we can compute that

$$E(T_G^n) = \sum_{(i_1, \dots, i_n) \in \{\pm 1, \dots, \pm N\}^n} E(T_{i_1} \dots T_{i_n}).$$

Therefore, by (5.4), we can obtain the following theorem.

Theorem 5.10. *Let T_G be the labeling operator of $\mathbb{G}$ on the graph Hilbert space H_G . Then*

(5.10)

$$E(T_G^n) = \sum_{w \in \mathcal{W}_n} R_w,$$

for all $n \in \mathbb{N}$, where $\mathcal{W}_n^{(m)}$ is defined by

$$\mathcal{W}_n^{(m)} \stackrel{\text{def}}{=} \left\{ w \in E(\widehat{G})^n \left| \begin{array}{l} w \in FP(\widehat{G}), \\ \omega_+(\omega(w)) = ((v, v), 0), \\ \text{for some } v \in V(G) \end{array} \right. \right\},$$

for all $n \in \mathbb{N}$. $\square$

The above theorem provides the D_G -valued free distributional data of the labeling operator T_G , in terms of D_G -valued moments. Similarly, we can have the following equivalent D_G -valued free distributional data of T_G , in terms of D_G -valued cumulants.

Theorem 5.11. *Let T_G be the labeling operator of $\mathbb{G}$ on H_G . Then the D_G -valued cumulants of T_G satisfy that: if n is odd, then $k_n(T_G, \dots, T_G) = 0_{D_G}$, and if n is even, then*

$$(5.11) \quad k_n \left(\underbrace{T_G, \dots, T_G}_{n\text{-times}} \right) = \sum_{(i_1, \dots, i_n) \in \{\pm 1, \dots, \pm N\}^n} \left(\sum_{w \in \mathcal{W}_n^{(c)}} \mu_w R_w \right),$$

where μ_w 's are defined in (5.7) and where $\mathcal{W}_n^{(c)}$ is defined by

$$(5.12) \quad \mathcal{W}_n^{(c)} \stackrel{\text{def}}{=} \bigcup_{(i_1, \dots, i_n) \in \{\pm 1, \dots, \pm N\}^n} \mathcal{W}_n^{(c)}(i_1, \dots, i_n), \text{ for all } n \in \mathbb{N}.$$

Here, $\mathcal{W}_n^{(c)}(i_1, \dots, i_n)$ is introduced in (5.9). $\square$

Recall that if an operator a of a B -valued W^* -probability space (A, E_B) , where $E_B : A \rightarrow B$ is a conditional expectation, is self-adjoint, then the B -valued free distribution σ_a of a and the (operator-valued) spectral measure χ_a of a are identical. Therefore, the formulae (5.10) and (5.11) also contain the spectral measure theoretical data of the labeling operator T_G (See [5] and [23]).

We will finish this section with the following corollary, which is the modification of (5.10). By the fact that all odd D_G -valued cumulants of T_G vanish, we can conclude that all odd D_G -valued moments of T_G vanish, too, by the Moebius Inversion. Indeed, we have

$$E(T_G^n) = \sum_{\pi \in NC(n)} k_{n:\pi}(T_G, \dots, T_G),$$

where $k_{n:\pi}(\bullet, \dots, \bullet)$ are the partition-depending D_G -valued cumulant. Notice that, if n is odd, then every noncrossing partition π of $NC(n)$ contains an odd block, having its odd size. Therefore, $k_{n:\pi}(T_G, \dots, T_G) = 0_{D_G}$, for all $\pi \in NC(n)$, and hence $E(T_G^n) = 0_{D_G}$, whenever n is odd. So, the formula (5.10) can be modified by:

Corollary 5.12. *Let T_G be the labeling operator of the given graph groupoid $\mathbb{G}$ on the graph Hilbert space H_G . Then*

(5.10)'

$$E(T_G^n) = \begin{cases} \sum_{w \in \mathcal{W}_n^{(m)}} R_w & \text{if } n \text{ is even} \\ 0_{D_G} & \text{if } n \text{ is odd,} \end{cases}$$

for all $n \in \mathbb{N}$. In particular, $\mathcal{W}_n^{(m)}$ is empty, whenever n is odd. $\square$

5.3. Graph-Theoretical Characterizations of $\mathcal{W}_{2n}^{(m)}$ and $\mathcal{W}_{2n}^{(c)}$.

As we have seen in (5.10) and (5.11), the D_G -valued free distributional data of the labeling operator T_G on the graph Hilbert space H_G is characterized by the sets $\{\mathcal{W}_{2n}^{(m)} : n \in \mathbb{N}\}$ and $\{\mathcal{W}_{2n}^{(c)} : n \in \mathbb{N}\}$, where

$$\mathcal{W}_{2n}^{(m)} = \left\{ w = e_1 \dots e_{2n} \left| \begin{array}{l} w \in E(\widehat{G})^{2n} \times FP(\widehat{G}), \\ \omega_+(\omega(w)) = ((v, v), 0), \\ \text{for some } v \in V(G) \end{array} \right. \right\}$$

and

$$\mathcal{W}_{2n}^{(c)} = \left\{ w = e_1 \dots e_{2n} \left| \begin{array}{l} w \in E(\widehat{G}) \times FP(\widehat{G}), \\ \exists e \in E(\widehat{G}) \text{ s.t., } w \text{ is the word} \\ \text{only in } \{e, e^{-1}\}, \text{ satisfying} \\ \omega_+(\omega(w)) = ((v, v), 0), \\ \text{for some } v \in V(G) \end{array} \right. \right\},$$

for all $n \in \mathbb{N}$. First, notice that both $\mathcal{W}_{2n}^{(m)}$ and $\mathcal{W}_{2n}^{(c)}$ are contained in $FP(\widehat{G}) \cap E(\widehat{G})^*$, where $E(\widehat{G})^* = \bigcup_{n=1}^{\infty} E(\widehat{G})^n (\supseteq FP(\widehat{G}))$ is the collection of all finite words in $E(\widehat{G})$, for all $n \in \mathbb{N}$, and $FP(\widehat{G})$ is the finite path set in the free semigroupoid $\mathbb{F}^+(\widehat{G})$ of the shadowed graph $\widehat{G}$ of the given graph G .

Now, we define the *loop-set*, $loop(\widehat{G})$ contained in the set $FP(\widehat{G}) \cap E(\widehat{G})^*$, by

$$loop(\widehat{G}) \stackrel{def}{=} \left\{ w \in E(\widehat{G})^* \left| \begin{array}{l} w \text{ is a loop} \\ \text{finite path} \\ \text{in } FP(\widehat{G}) \end{array} \right. \right\} \subseteq FP(\widehat{G}).$$

More precisely, we can define it by

$$loop(\widehat{G}) \stackrel{def}{=} \bigcup_{v \in V(G)} \left(loop_v(\widehat{G}) \right),$$

where

$$\text{loop}_v(\widehat{G}) \stackrel{\text{def}}{=} \{w \in FP(\widehat{G}) : w = v w v\}.$$

i.e., the subset $\text{loop}(\widehat{G})$ of $FP(\widehat{G})$ is decomposed by $\text{loop}_v(\widehat{G})$, for all $v \in V(G)$. Clearly, if a vertex v of G has no incident loop finite path, then $\text{loop}_v(\widehat{G})$ is empty. However, since $\text{loop}_v(\widehat{G})$ are contained in $\mathbb{F}^+(\widehat{G})$ (not in $\mathbb{G}$), and since our graph G (and $\widehat{G}$) is connected, whenever there is at least one edge e such that either $e = v$ or $e = e v$, the set $\text{loop}_v(\widehat{G})$ is nonempty. For instance, if $e = ve$ in $E(\widehat{G})$, then $ee^{-1} \in \text{loop}_v(\widehat{G})$.

By the definition of $\text{loop}(\widehat{G})$, we can re-construct the set $\mathcal{W}_{2n}^{(m)}$:

$$\mathcal{W}_{2n}^{(m)} = \left\{ w \in E(\widehat{G})^{2n} \left| \begin{array}{l} w \in \text{loop}(\widehat{G}), \text{ satisfying} \\ \omega_+(\omega(w)) = ((v, v), 0), \\ \text{for some } v \in V(G) \end{array} \right. \right\},$$

for all $n \in \mathbb{N}$.

Now, recall the reduction (RR) on the free semigroupoid $\mathbb{F}^+(\widehat{G})$. We give the reduction (RR), as the following quotient map $\delta : \mathbb{F}^+(\widehat{G}) \rightarrow \mathbb{G}$ (and hence surjective), defined by

$$\delta(W) = w \in \mathbb{G}, \text{ for all } W \in \mathbb{F}^+(\widehat{G}),$$

satisfying that

$$\delta(W W^{-1}) = v_1 \text{ and } \delta(W^{-1} W) = v_2,$$

for all $W = v_1 W v_2 \in \mathbb{F}^+(\widehat{G})$, with $v_1, v_2 \in V(\widehat{G}) = V(G)$. By [62], we can check that, indeed, the map $\delta : \mathbb{F}^+(\widehat{G}) \rightarrow \mathbb{G}$ is the well-defined quotient map. Equivalently,

$$\mathbb{G} \stackrel{\text{Groupoid}}{=} \mathbb{F}^+(\widehat{G}) / \delta(\mathbb{F}^+(\widehat{G})).$$

We say that δ is the *reduction (quotient) map of $\mathbb{F}^+(\widehat{G})$ onto $\mathbb{G}$* . By help of this map δ , we can re-construct the set $\mathcal{W}_{2n}^{(m)}$ by

$$\mathcal{W}_{2n}^{(m)} = \left\{ w \in E(\widehat{G})^{2n} \cap \text{loop}(\widehat{G}) \mid \delta(w) \in V(\widehat{G}) \right\},$$

for all $n \in \mathbb{N}$. Thus the set $\mathcal{W}_{2n}^{(n)}$ is characterized, graph-theoretically, as follows.

Proposition 5.13. *For $n \in \mathbb{N}$, we have that*

$$\mathcal{W}_{2n}^{(m)} = \left\{ w \in E(\widehat{G})^{2n} \cap \text{loop}(\widehat{G}) \mid \delta(w) \in V(\widehat{G}) \right\}.$$

□

Now, we define the subset $\mathcal{D}_{\text{loop}}(\widehat{G})$ of $\text{loop}(\widehat{G})$ in $FP(\widehat{G}) \subset \mathbb{F}^+(\widehat{G})$ by

$$\mathcal{D}_{\text{loop}}(\widehat{G}) \stackrel{\text{def}}{=} \left\{ w \in \text{loop}(\widehat{G}) \mid \begin{array}{l} \exists e \in E(\widehat{G}) \text{ s.t.} \\ w \text{ is a word} \\ \text{only in } \{e, e^{-1}\} \end{array} \right\}.$$

Then we can re-construct $\mathcal{W}_{2n}^{(c)}$ by

$$\mathcal{W}_{2n}^{(c)} = \left\{ w \in E(\widehat{G})^{2n} \mid \begin{array}{l} w \in \mathcal{D}_{\text{loop}}(\widehat{G}), \text{ satisfying} \\ \omega_+(\omega(w)) = ((v, v), 0), \\ \text{for some } v \in V(G) \end{array} \right\},$$

for all $n \in \mathbb{N}$. Therefore, we obtain the following proposition.

Proposition 5.14. *For any $n \in \mathbb{N}$, we have that*

$$\mathcal{W}_{2n}^{(c)} = \left\{ w \in E(\widehat{G})^{2n} \mid \begin{array}{l} w \in \mathcal{D}_{\text{loop}}(\widehat{G}), \\ \text{satisfying} \\ \delta(w) \in V(\widehat{G}) \end{array} \right\}.$$

□

The above two propositions provide the combinatorial characterization of the sets $\mathcal{W}_{2n}^{(m)}$ and $\mathcal{W}_{2n}^{(c)}$, for $n \in \mathbb{N}$. It means that we can verify the D_G -valued moments and cumulants of the labeling operator T_G of $\mathbb{G}$ on the graph Hilbert space H_G , simply by checking the shadowed graph $\widehat{G}$ of G .

6. EXAMPLES

In this section, we give several examples of the labeling operators of the labeled graph groupoids. In Section 4, we showed that the D_G -valued free distributional data of the labeling operator T_G of a (labeled) graph groupoid $\mathbb{G}$ on H_G is characterized by the certain subsets $\mathcal{W}_n^{(m)}$ and $\mathcal{W}_{2n}^{(c)}$ of the (non-reduced) finite path set $FP(\widehat{G})$ of the free semigroupoid $\mathbb{F}^+(\widehat{G})$, where $\widehat{G}$ is the shadowed graph of G , for all $n \in \mathbb{N}$. The D_G -valued free distributional data of the labeling operator T_G of $\mathbb{G}$ is represented by the D_G -valued moments $\{E(T_G^n)\}_{n \in \mathbb{N}}$, equivalently, by the

D_G -valued cumulants $\{k_n(T_G, \dots, T_G)\}_{n \in \mathbb{N}}$, where $E : M_G \rightarrow D_G$ is the canonical conditional expectation;

$$E(T_G^{2n}) = \sum_{w \in \mathcal{W}_{2n}^{(m)}} R_w, \text{ for all } n \in \mathbb{N},$$

and

$$k_{2n} \left(\underbrace{T_G, \dots, T_G}_{2n\text{-times}} \right) = \sum_{w \in \mathcal{W}_{2n}^{(c)}} (\mu_w R_w), \text{ for all } n \in \mathbb{N},$$

with

$$k_n(T_G, \dots, T_G) = 0_{D_G} = E(T_G^n),$$

for all odd n .

Example 6.1. (Fractaloid Cases) Let G be a canonical weighted graph with its labeled graph groupoid $\mathbb{G}$, and assume that $\mathbb{G}$ is a fractaloid. i.e., the automata actions $\{\mathcal{A}_w : w \in \mathbb{F}^+(\widehat{G})\}$ of the G -automaton $\mathcal{A}_G$ acts fully on the $\mathcal{A}_G$ -tree $\mathcal{T}_{2N}$, where $\mathcal{T}_{2N}$ is the $(2N)$ -regular tree, where $N = \max\{\deg_{out}(v) : v \in V(G)\}$ in $\mathbb{N}$. Now, let T_G be the labeling operator of the fractaloid $\mathbb{G}$ on H_G . Consider now the subset $\mathcal{W}_{2n}^{(m)}$ of the finite path set $FP(\widehat{G})$ of the shadowed graph $\widehat{G}$ of G , for all $n \in \mathbb{N}$. Recall that

$$\mathcal{W}_{2n}^{(m)} = \left\{ w \in E(\widehat{G})^{2n} \mid \begin{array}{l} \omega_+(\omega(w)) = ((v, v), 0) \\ \text{for some } v \in V(G) \end{array} \right\},$$

contained in $\text{loop}(\widehat{G}) \subseteq FP(\widehat{G})$. Moreover, the D_G -valued moments $E(T_G^{2n})$ is determined by

$$E(T_G^{2n}) = \sum_{w \in \mathcal{W}_{2n}^{(m)}} R_w, \text{ for all } n \in \mathbb{N}.$$

We can verify that, since $\mathbb{G}$ is a fractaloid, for any $n \in \mathbb{N}$, we have

$$(6.1) \quad \sum_{w \in \mathcal{W}_{2n}^{(m)}} R_w = \sum_{(l_{i_1}, \dots, l_{i_{2n}}) \in (\pm X_0^*)^{2n}, \theta(l_{i_1} \dots l_{i_{2n}}) = 0} R_{e_1 \dots e_{2n}},$$

where $e_1 \dots e_{2n} \in E(\widehat{G})^{2n}$, with $(\tilde{p}r \circ \omega)(e_j) = l_{i_j}$, for all $j = 1, \dots, 2n$. Then, again, since $\mathbb{G}$ is a fractaloid, there are $|V(G)|$ -many words $w_1, \dots, w_{|V(G)|}$, in $E(\widehat{G})^{2n}$ (possibly, $|V(G)| = \infty$) satisfying that $w_k = e_{k_1} \dots e_{k_{2n}}$ such that $(\tilde{p}r \circ \omega)(e_{k_q}) = l_{i_q}$, for $q = 1, \dots, 2n$. Since $\sum_{v \in V(G)} R_v$ is the identity element $1_{M_G} = 1_{D_G}$ of the right graph von Neumann algebra M_G , the formula (6.1) is identical to

$$(6.2)$$

$$\sum_{v \in V(G)} \left(\left| \sum_{2n}^{(N)} \right| \cdot R_v \right) = |\mathcal{L}_N^o(2n)| \left(\sum_{v \in V(G)} R_v \right) = |\mathcal{L}_N^o(2n)| \cdot 1_{D_G},$$

where $\mathcal{L}_N^o(2n)$ are introduced in Section 3.1, for all $n \in \mathbb{N}$. In [17], we showed that, the cardinalities $|\mathcal{L}_N^o(2n)|$ are gotten recursively by the entries of the famous Pascal's triangle (Also, see [41]).

This example shows that if $\mathbb{G}$ is a fractaloids, then the amalgamated (or the operator-valued) free distributional data of the labeling operator is completely determined by the scalar-values, which are the cardinalities of certain subsets of the lattice path set $\mathcal{L}_N$.

Remember that, in [17], we conjectured that the only “finite” connected directed graphs inducing fractaloids are one-flow circulant graphs and one-vertex-multi-loop-edge graphs and the finite iterated glued graphs of them, in the sense of [16]. This conjecture is still open. If it is positive, then the finite fractaloids are completely characterized, and hence we could obtain the complete analysis of their labeling operators in terms of their amalgamated free distributional data.

Example 6.2. Let G be a directed graph with

$$V(G) = \{v_1, v_2, v_3\}$$

and

$$E(G) = \{e_{12:1}, e_{12:2}, e_{13:1}, e_{22:1}\},$$

where $e_{ij:k}$ means the k -th edge connecting the vertex v_i to the vertex v_j . Then we can have the labeling set

$$X = \{l_1 = \overrightarrow{(1, e)}, l_2 = \overrightarrow{(1, e^2)}\} \subset \mathcal{L}_2$$

since

$$\max\{\deg_{out}(v_k) : k = 1, 2, 3\} = 2 = \deg_{out}(v_1).$$

Then the weighting process $\omega : \mathbb{F}^+(\widehat{G}) \rightarrow V(G)^2 \times \pm X_0^* = \mathcal{X}_0^*$ is completely determined by the indices of the edges:

$$\omega(e_{ij:k}) = ((v_i, v_j), l_k).$$

and

$$\omega(e_{ij:k}^{-1}) = ((v_j, v_i), -l_k).$$

By definition, we can have the k -th labeling operators, for $k = \pm 1, \pm 2$, as elements in the right graph von Neumann algebra M_G as follows:

$$T_1 = R_{e_{12:1}} + R_{e_{13:1}} + R_{e_{22:1}} \text{ and } T_{-1} = T_1^*$$

and

$$T_2 = R_{e_{12:2}}, \text{ and } T_{-2} = T_2^* = R_{e_{12:2}^{-1}}.$$

and hence

$$T_G = (T_1 + T_{-1}) + (T_2 + T_{-2}).$$

Then we can have that $E(T_G^k) = 0_{D_G}$, whenever k is odd, and

$$E(T_G^{2n}) = \sum_{w \in \mathcal{W}_{2n}^{(m)}} R_w, \text{ for all } n \in \mathbb{N},$$

by Section 5.2. In our case, we have that

$$\mathcal{W}_2^{(m)} = \left\{ \begin{array}{ccc} e_{12:1}e_{12:1}^{-1}, & e_{12:1}^{-1}e_{12:1}, & e_{12:2}e_{12:2}^{-1}, \\ e_{12:2}^{-1}e_{12:2}, & e_{13:1}e_{13:1}^{-1}, & e_{13:1}^{-1}e_{13:1} \end{array} \right\},$$

by Section 5.3. So, we can have that

$$\begin{aligned} E(T_G^2) &= R_{e_{12:1}e_{12:1}^{-1}} + R_{e_{12:1}^{-1}e_{12:1}} + R_{e_{12:2}e_{12:2}^{-1}} \\ &\quad + R_{e_{12:2}^{-1}e_{12:2}} + R_{e_{13:1}e_{13:1}^{-1}} + R_{e_{13:1}^{-1}e_{13:1}} \\ &= R_{v_1} + R_{v_2} + R_{v_1} + R_{v_2} + R_{v_1} + R_{v_3} \\ &= 3 R_{v_1} + 2 R_{v_2} + R_{v_3}. \end{aligned}$$

The above computation $E(T_G^2)$ of T_G says that the given graph G does not generate a fractaloid.

7. !!!!!!!!!!!!!!!PUT PICTURES!!!!!!!!!!!!!!!!!!!!!! $G\mathcal{G}\omega\varphi\psi\tilde{p}r$!!!!!!!!!!!!!!!!!!!!!!!!!!!!!!

References

- [1] A. G. Myasnikov and V. Shpilrain (editors), Group Theory, Statistics and Cryptography, Contemporary Math, 360, (2003) AMS.
- [2] A. Gibbons and L. Novak, Hybrid Graph Theory and Network Analysis, ISBN: 0-521-46117-0, (1999) Cambridge Univ. Press.
- [3] B. Solel, You can see the arrows in a Quiver Operator Algebras, (2000), preprint.
- [4] C. W. Marshall, Applied Graph Theory, ISBN: 0-471-57300-0 (1971) John Wiley & Sons
- [5] D. Voiculescu, K. Dykemma and A. Nica, Free Random Variables, CRM Monograph Series Vol 1 (1992).
- [6] D.W. Kribs and M.T. Jury, Ideal Structure in Free Semigroupoid Algebras from Directed Graphs, preprint.
- [7] D.W. Kribs, Quantum Causal Histories and the Directed Graph Operator Framework, arXiv:math.OA/0501087v1 (2005), Preprint.
- [8] F. Balacheff, Volume Entropy, Systole and Stable Norm on Graphs, arXiv:math.MG/0411578v1, (2004) Preprint.
- [9] G. C. Bell, Growth of the Asymptotic Dimension Function for Groups, (2005) Preprint.
- [10] I. Cho, Graph von Neumann algebras, ACTA. Appl. Math, 95, (2007) 95 - 135.
- [11] I. Cho, Characterization of Free Blocks of a right graph von Neumann algebra, Compl. An. & Op. theo (2007) To be Appeared.
- [12] I. Cho, Direct Producted W^* -Probability Spaces and Corresponding Free Stochastic Integration, B. of KMS, 44, No. 1, (2007), 131 - 150.
- [13] I. Cho, Vertex-Compressed Algebras of a Graph von Neumann Algebra, (2007) Submitted to ACTA. Appl. Math.
- [14] I. Cho, Group-Freeness and Certain Amalgamated Freeness, J. of KMS, 45, no. 3, (2008) 597 - 609.
- [15] I. Cho and P. E. T. Jorgensen, C^* -Algebras Generated by Partial Isometries, JAMC, (2008) To Appear.
- [16] I. Cho and P. E. T. Jorgensen, C^* -Subalgebras Generated by Partial Isometries, JMP, (2008) To Appear.
- [17] I. Cho and P. E. T. Jorgensen, Applications of Automata and Graphs: Labeling-Operators in Hilbert Space I, (2008) Submitted to ACTA Appl. Math., Special Issues.
- [18] I. Raeburn, Graph Algebras, CBMS no 3, AMS (2005).
- [19] P. D. Mitchener, C^* -Categories, Groupoid Actions, Equivalent KK-Theory, and the Baum-Connes Conjecture, arXiv:math.KT/0204291v1, (2005), Preprint.
- [20] R. Scapellato and J. Lauri, Topics in Graph Automorphisms and Reconstruction, London Math. Soc., Student Text 54, (2003) Cambridge Univ. Press.
- [21] R. Exel, A new Look at the Crossed-Product of a C^* -algebra by a Semigroup of Endomorphisms, (2005) Preprint.
- [22] R. Gliman, V. Shpilrain and A. G. Myasnikov (editors), Computational and Statistical Group Theory, Contemporary Math, 298, (2001) AMS.

- [23] R. Speicher, Combinatorial Theory of the Free Product with Amalgamation and Operator-Valued Free Probability Theory, AMS Mem, Vol 132 , Num 627 , (1998).
- [24] S. H. Weintraub, Representation Theory of Finite Groups: Algebra and Arithmetic, Grad. Studies in Math, vo. 59, (2003) AMS.
- [25] V. Vega, Finite Directed Graphs and W^* -Correspondences, (2007) Ph. D thesis, Univ. of Iowa.
- [26] W. Dicks and E. Ventura, The Group Fixed by a Family of Injective Endomorphisms of a Free Group, Contemp. Math 195, AMS.
- [27] F. Radulescu, Random Matrices, Amalgamated Free Products and Subfactors of the von Neumann Algebra of a Free Group, of Noninteger Index, Invent. Math., 115, (1994) 347 - 389.
- [28] D. A. Lind, Entropies of Automorphisms of a Topological Markov Shift, Proc. AMS, vo 99, no 3, (1987) 589 - 595.
- [29] D. A. Lind and B. Marcus, An Introduction to Symbolic Dynamics and Coding, (1995) Cambridge Univ. Press.
- [30] D. A. Lind and S. Tuncel, A Spanning Tree Invariant for Markov Shifts, IMA Vol. Math. Appl., vo 123, (2001), 487 - 497.
- [31] D. A. Lind and K. Schmidt, Symbolic and Algebraic Dynamical Systems, Handbook of Dynamical System, Vol. 1A, (2002) 765 - 812.
- [32] R. V. Kadison and J. R. Ringrose, Fundamentals of the Theory of Operator Algebra, Grad. Stud. Math., vo. 15, (1997) AMS.
- [33] D. E. Dutkay and P. E. T. Jorgensen, Iterated Function Systems, Ruelle Operators and Invariant Projective Measures, arXiv:math.DS/0501077/v3, (2005) Preprint.
- [34] P. E. T. Jorgensen, Use of Operator Algebras in the Analysis of Measures from Wavelets and Iterated Function Systems, (2005) Preprint.
- [35] D. Guido, T. Isola and M. L. Lapidus, A Trace on Fractal Graphs and the Ihara Zeta Function, arXiv:math.OA/0608060v1, (2006) Preprint.
- [36] P. Potgieter, Nonstandard Analysis, Fractal Properties and Brownian Motion, arXiv:math.FA/0701649v1, (2007) Preprint.
- [37] L. Bartholdi, R. Grigorchuk, and V. Nekrashevych, From Fractal Groups to Fractal Sets, arXiv:math.GR/0202001v4, (2002) Preprint.
- [38] I. Cho, The Moments of Certain Perturbed Operators of the Radial Operator of the Free Group Factor $L(F_N)$, JAA, 5, no. 3, (2007) 137 - 165.
- [39] I. Cho and P. E. T. Jorgensen, C^* -Subalgebras Generated by Partial Isometries in $B(H)$, JMP (2008) To Appear.
- [40] S. Thompson and I. Cho, Powers of Mutinomials in Commutative Algebras, (2008) (Undergraduate Research) Submitted to PMEJ.
- [41] S. Thompson, C. M. Mendoza, and A. J. Kwiatkowski, and I. Cho, Lattice Paths Satisfying the Axis Property, (2008) (Undergraduate Research) Preprint.
- [42] I. Cho, Labeling Operators of Graph Groupoids, (2008) Preprint.
- [43] R. T. Powers, Heisenberg Model and a Random Walk on the Permutation Group, Lett. Math. Phys., 1, no. 2, (1975) 125 - 130.

- [44] R. T. Powers, Resistance Inequalities for KMS -states of the isotropic Heisenberg Model, *Comm. Math. Phys.*, 51, no. 2, (1976) 151 - 156.
- [45] R. T. Powers, Resistance Inequalities for the Isotropic Heisenberg Ferromagnet, *JMP*, 17, no. 10, (1976) 1910 - 1918.
- [46] E. P. Wigner, Characteristic Vectors of Bordered Matrices with Infinite Dimensions, *Ann. of Math.* (2), 62, (1955) 548 - 564.
- [47] D. Voiculescu, Symmetries of Some Reduced Free Product C^* -Algebras, *Lect. Notes in Math.*, 1132, Springer, (1985) 556 - 588.
- [48] T. Shirai, The Spectrum of Infinite Regular Line Graphs, *Trans. AMS.*, 352, no 1., (2000) 115 - 132.
- [49] J. Kigami, R. S. Strichartz, and K. C. Walker, Constructing a Laplacian on the Diamond Fractal, *Experiment. Math.*, 10, no. 3, (2001) 437 - 448.
- [50] I. V. Kucherenko, On the Structurization of a Class of Reversible Cellular Automata, *Diskret. Mat.*, 19, no. 3, (2007) 102 - 121.
- [51] J. L. Schiff, Cellular Automata, Discrete View of the World, Wiley-Interscience Series in Disc. Math. & Optimazation, ISBN: 978-0-470-16879-0, (2008) John Wiley & Sons Press.
- [52] P. E. T. Jorgensen, and M. Song, Entropy Encoding, Hilbert Spaces, and Kahunen-Loeve Transforms, *JMP*, 48, no. 10, (2007)
- [53] P. E. T. Jorgensen, L. M. Schmitt, and R. F. Werner, q -Canonical Commutation Relations and Stability of the Cuntz Algebra, *Pac. J. of Math.*, 165, no. 1, (1994) 131 - 151.
- [54] A. Gill, Introduction to the Theory of Finite-State Machines, MR0209083 (34\#8891), (1962) McGraw-Hill Book Co.
- [55] J. E. Hopcroft, and J. D. Ullman, Introduction to Automata Theory, Language, and Computation, ISBN: 0-201-02988-X, (1979) Addison-Wesley Publication Co.
- [56] M. Fannes, B. Nachtergaele, and R. F. Werner, Ground States of VSB -Models on Cayley Trees, *J. of Statist. Phys.*, 66, (1992) 939 - 973.
- [57] M. Fannes, B. Nachtergaele, and R. F. Werner, Finitely Correlated States on Quantum Spin Chains, *Comm. Math. Phys.*, 144, no. 3, (1992) 443 - 490.
- [58] M. Fannes, B. Nachtergaele, and R. F. Werner, Entropy Estimates for Finitely Correlated States, *Ann. Inst. H. Poincare. Phys. Theor.*, 57, no 3, (1992) 259 - 277.
- [59] M. Fannes, B. Nachtergaele, and R. F. Werner, Finitely Correlated Pure States, *J. of Funct. Anal.*, 120, no 2, (1994) 511 - 534.
- [60] J. Renault, A Groupoid Approach to C^* -Algebras, *Lect. Notes in Math.*, 793, ISBN: 3-540-09977-8, (1980) Springer.
- [61] S. Sakai, C^* -Algebras and W^* -Algebras, MR number: MR0442701, (1971) Springer-Verlag.
- [62] I. Cho, Measures on Graphs and Groupoid Measures, *Compl. An. Oper. Theor.*, 2, (2008) 1 - 28.
- [63] B. Fuglede, and R. V. Kadison, On Determinants and a Property of the Trace in Finite Factors, *Proc. Nat. Acad. Sci., U. S. A.*, 37, (1951) 425 - 431.

- [64] D. Aldous, and R. Lyons, Processes on Unimodular Random Networks, Elec. J. of Probab., 12, no 54, (2007) 1454 - 1508.
- [65] O. Gühhne, G. Tóth, P. Hyllus, and H. J. Briegel, Bell Inequalities for Graphs States, Phys. Rev. Lett., 95, no 12, (2005)
- [66] R. V. Kadison, and J. R. Ringrose, Fundamentals of the Theory of Operator Algebras, vol II, Grad. Stud. in Math., 16, ISBN: 0-8218-0820-6, (1997) AMS.

SAINT AMBROSE UNIV., DEP. OF MATH., 421 AMBROSE HALL, 518 W. LOCUST ST., DAVENPORT, IOWA, 52803, U. S. A. / UNIVERSITY OF IOWA, DEP. OF MATH., 14 MCLEAN HALL, IOWA CITY, IOWA, 52242-1419, U. S. A.

E-mail address: `chowoo@sau.edu`, `jorgense@math.uiowa.edu`